



MONTHLY ANALYSIS OF PM₁₀ IN AMBIENT AIR OF KLANG VALLEY, MALAYSIA

(Analisis PM₁₀ Bulanan di dalam Udara di Lembah Klang, Malaysia)

Mohd Asrul Jamalani¹, Ahmad Makmom Abdullah^{1,2*}, Azman Azid^{3,4}, Mohammad Firuz Ramli²,
Mohd Rafee Baharudin⁵, Mahmud Mohammed Bose¹, Rashieda Elawad Elhadi¹,
Khaleed Ali Ahmed Ben Youssef¹, Azadeh Gnadimzadeh¹, Danladi Yusuf Gumel¹

¹*Air Quality and Ecophysiology Laboratory, Faculty of Environmental Studies*

²*Department of Environmental Sciences, Faculty of Environmental Studies*

Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia

³*UniSZA Science and Medicine Foundation Centre,*

Universiti Sultan Zainal Abidin, Gong Badak Campus, 21300 Kuala Nerus, Terengganu, Malaysia

⁴*Faculty Bioresources and Food Industry,*

Universiti Sultan Zainal Abidin, Tembila Campus, 22200 Besut, Terengganu, Malaysia

⁵*Department of Community Health, Faculty of Medicine and Health Sciences,*

Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia

*Corresponding author: amakmom@upm.edu.my

Received: 14 April 2015; Accepted: 3 August 2016

Abstract

The urbanization in Klang Valley, Peninsular Malaysia over the last decades has induce the atmospheric pollution's risk resulted to negative impact on the environment. The aims of this paper are to identify the spatial-temporal relationship of particulate matter (PM₁₀), to determine the characteristic of each location and to classify hierarchical of the location in relation to their impact on PM₁₀ concentration in Klang Valley. The Spearman correlation test indicate that there was strong significant relationship between all the locations (> 0.7 ; $p < 0.001$) and moderate relationship between Petaling Jaya-Kajang and Kajang-Shah Alam (< 0.7 ; $p < 0.001$). The principal component analysis (PCA) identifies all four locations have been affected by PM₁₀ which were determined as one of the pollutant that deteriorated the air quality. Cluster analysis (CA) has classified the PM₁₀ pattern into three (3) different classes; Class 1 (Klang), Class 2 (Petaling Jaya and Kajang) and Class 3 (Shah Alam) based on location. Further analysis of CA would be able to classify the PM₁₀ classes into groups depending on their dissimilarities characteristic. Thus, possible period of extreme air quality degradation could be identified. Therefore, statistical and envirometric techniques have proved the impact of the various location on increasing concentration of PM₁₀.

Keywords: particulate matter, Spearman correlation test, principal component analysis, cluster analysis

Abstrak

Proses pembandaran di Lembah Klang, Semenanjung Malaysia sedekad lalu telah mendorong kepada risiko pencemaran atmosfera yang memberi impak negatif kepada alam sekitar. Kajian ini dilakukan bertujuan untuk mengenalpasti hubungkait antara ruang dan tempoh bagi partikel terampai (PM₁₀), menentukan ciri – ciri setiap lokasi dan menentukan pengkelasan hirarki lokasi berhubungan dengan impak kepekatan PM₁₀ di Lembah Klang. Ujian korelasi Spearman menunjukkan hubungkait yang kuat antara semua lokasi (> 0.7 ; $p < 0.001$) dan hubungan

yang sederhana antara Petaling Jaya-Kajang dan Kajang-Shah Alam (< 0.7 ; $p < 0.001$). Analisis komponen utama (PCA) menentukan semua empat lokasi yang telah terjejas dengan PM₁₀ iaitu antara bahan pencemar yang menjejaskan kualiti udara. Analisis kluster (CA) mengelaskan pola PM₁₀ kepada tiga (3) kelas berlainan; Kelas 1 (Klang), Kelas 2 (Petaling Jaya dan Kajang) serta Kelas 3 (Shah Alam) berdasarkan lokasi. Analisis lanjutan CA membolehkan pengelasan kelas PM₁₀ kepada kumpulan bergantung kepada ketidaksamaan ciri. Justeru, kemungkinan tempoh kemerosotan kualiti udara yang melampau dapat dikenalpasti. Oleh itu, teknik statistik dan envirometrik telah membuktikan impak pelbagai lokasi terhadap peningkatan kepekatan PM₁₀.

Kata kunci: partikel terampai, ujian korelasi Spearman, analisis komponen utama, analisis kluster

Introduction

Particulate matter is one of the aerosol particles in the atmosphere [1, 2] which has an aerodynamic diameter of less than 10 μm and well known as PM₁₀. PM₁₀ has been discovered as major air pollutant in Southeast Asia including Klang Valley, Malaysia [3, 4, 5]. A long term study over 5 decades regarding to air pollution brings up an implication not only to human health but also to the environment [6, 7]. Particulate matter specifically PM₁₀ affects human health via inhalation due to its smaller in size and the ability to reach and settle in human respiratory tract which could induce chronic pulmonary disease and asthma [8, 9].

Asia region encounter a major problem due to air particulate matter pollution with the annual average value of total suspended particulate (TSP) which exceeding 300 $\mu\text{g}\text{m}^{-3}$ [10]. In Malaysia without the haze issue, the level of PM₁₀ is mostly influence by the vehicular and industrial emission [4]. Therefore, this study will define the PM₁₀ level in Klang Valley region under normal condition without severe haze issue as the main focus.

Multivariate analysis has been chosen as the final statistical method to analyze, classify and interpret huge number of datasets and has become the most applicable in various field of study recently [11 - 19]. These types of analysis include the application of Principal Component Analysis (PCA) and Hierarchical Agglomerative Cluster Analysis (HACA). The PCA was applying to identify the most significant parameter which relates to spatial and temporal variation [11 - 17, 20, 21, 22]. Meanwhile, HACA was applying to group large data into cluster with differing characteristic between the groups but similar characteristic within the group [23]. Therefore, the aim of this study is to identify the spatial-temporal relationship of PM₁₀, to determine the characteristic of each station and to classify hierarchical of the station that give an impact to PM₁₀ concentration in Klang Valley.

Materials and Methods

Background of sampling location

Klang Valley region involves several districts in Selangor and is located in central part of west coast Peninsular Malaysia by the Strait of Malacca to west [24, 25]. Alam Sekitar Malaysia Sdn. Bhd. (ASMA) was appointed by Malaysian Department of Environment (DOE) in establishing, operating and maintaining the continuous air quality monitoring stations. All the stations (S1, S3 and S4) were located within residential area except for Petaling Jaya (S2) which located within industrial area. All the stations within Klang Valley region were affected by heavy traffic which consequently affected by vehicles emissions and the details were shown in the Table 1. The Klang Valley Region which is situated in the central part of Selangor state is illustrated in Figure 1.

The air quality data

The daily-recorded air quality data of PM₁₀ at the selected stations within the year of 2000 – 2009 were obtained from Malaysian Department of Environment (DOE). A total of 480 observation data of PM₁₀ (12 observations x 4 stations x 10 years) were involved in this study.

Data cleaning

Data treatment technique has been used to treat the missing data in order to obtain a better continuous air quality monitoring data. The missing data in this study has been treated with the method of mean substitution. All the missing data are replacing with the value of mean available neighboring data [27]. Furthermore, other studies stated

that mean substitution method is better and more accurate rather than eliminating the missing value with list wise and pairwise deletion method [28, 29, 30].

Table 1. Continuous air quality monitoring stations within Klang Valley region

Station ID	Air Monitoring Station	Representative Station ID	Representative Name	Coordinates		Background
				Latitude (N)	Longitude (E)	
CA0011	SM(P) Raja Zarina, Klang	S1	Klang	3.0100°	101.4085°	Residential
CA0016	Sek. Keb. Seri Petaling, Petaling Jaya	S2	Petaling Jaya	3.1092°	101.6387°	Industry
CA0023	Country Height, Kajang	S3	Kajang	2.9939°	101.7417°	Residential
CA0025	Sek. Keb. TTDI, ShahAlam	S4	Shah Alam	3.0773°	101.5112°	Residential

Source: Department of Environment [26]




Figure 1. Map of Selangor district which include Klang Valley region

Data analysis

Several types of statistical approach include descriptive analysis, Spearman correlation analysis, principal component analysis (PCA) and cluster analysis (CA) via XLSTAT software has been used in achieving the objective. XLSTAT software has been used as a tool because of its flexibility, multidimensionality and ability to synthesize complex data sets [15]. Any missing value had been treated using mean substitution method.

Spearman correlation test

Spearman correlation test is used for determination of relationship between two variables and also known as Spearman Rho test [31, 32, 33]. This test is one of the non-parametric approaches which is suitable for not normally distributed data and would prefer monotonic graph instead of linear graph. Monotonic relationship shows that the score of variable increases with decreasing the other variables' score either in similar or different rate. The correlation values are in the range between -1 and +1 where the value only shows the strength of correlation between two variables without showing cause and consequence between those two variables [34]. However, the influence and the impact of each other variables can be described. Mathematical formula for calculating Spearman correlation is as follows;

$$r = 1 - \left(\frac{6 \sum p^2}{N(N^2 - 1)} \right) \quad (1)$$

where p^2 is define as square root of the variable, and N is a sample size

Principal component analysis (PCA)

Principal component analysis is a procedure for identifying, reducing and arranging items into groups depending on dependent variables and the strength of correlation between those items [12, 15, 16, 35 – 39]. Linear combination of original data set is being generated by the ability of PCA in reducing large amount of data into new sets of variables where the number of principal components is not more than number of original variables [37]. Identification and observation of variation's source taking place after reduction of data set and generally written in the following mathematical equation 2;

$$PC_i = l_{1i} X_1 + l_{2i} X_2 + \dots + l_{ni} X_n \quad (2)$$

where PC_i is define as i^{th} principal component, l_{ji} is define as variable loading and X_j is define as observed variable

Cluster analysis (CA)

CA is an unsupervised pattern recognition identification method, used to split a large group into smaller ones [12] based on homogeneity data. The homogeneous sub-groups will be obtained within the population and gather them into clusters based on similarity of the data [38, 40, 41]. In this study, CA was used for clustering data with the similarities in a group. CA is employed on the normal distribution dataset through the Ward's method by means of Euclidean distances, as a measure of the relationship [11, 12, 16]. The outcome of this method will be demonstrated in a dendrogram form.

Results and Discussion**Statistical analysis of PM₁₀**

The monthly values of PM₁₀ data in Klang Valley, Selangor have been analyzed and summarized in box plots. The PM₁₀ distribution in the air is higher in Klang and Shah Alam compare to Petaling Jaya and Kajang with the range of 154.65 $\mu\text{g}/\text{m}^3$ and 104.19 $\mu\text{g}/\text{m}^3$ respectively. The mean of PM₁₀ is also higher in Klang and Shah Alam compare to the other two stations with the value of 79.97 $\mu\text{g}/\text{m}^3$ and 63.32 $\mu\text{g}/\text{m}^3$ respectively. Major activities and heavy populated with residential and industrial activities are the main contributors to the higher PM₁₀ value in Klang Valley region (Klang and Shah Alam). The details of descriptive statistic of PM₁₀ distributions in Klang Valley region for 2000 to 2009 is shown in Table 2 and expressed by box plot in Figure 2.

Determination of air quality relationship

The monthly PM₁₀ data in Klang Valley region for 10 years period (2000 to 2009) showed the result is not normally distributed. Parametric test has been used for transformation process via log method. However, the negative result still not normally distributed form. Therefore, the non-parametric test was chosen for the PM₁₀ data instead of parametric test. Thus, the Spearman correlation was chosen as the non-parametric approach for determining the relationship between two variables.

Table 2. Descriptive statistic of monthly PM₁₀ distributions in Klang Valley region for year 2000-2009

Station	Obs	Min	Max	Range	Q1	Median	Q3	Mean	Var (n)	Std. dev. (n)
Klang	120	39.25	193.90	154.65	55.14	68.39	95.02	79.97	1149.72	33.91
Petaling Jaya	120	26.82	116.97	90.15	45.34	54.06	67.45	58.28	316.60	17.79
Kajang	120	31.01	124.48	93.48	42.30	49.87	61.73	54.03	283.27	16.83
Shah Alam	120	29.67	133.86	104.19	44.71	58.71	80.07	63.32	535.08	23.13




Figure 2. Box Plot of PM₁₀ distributions in Klang Valley region for 10 years period (2000 – 2009)

Table 3 showed that all the stations have significant relationship. However, only Klang-Petaling Jaya, Klang-Kajang, Klang-Shah Alam and Petaling Jaya-Shah Alam prove for strong correlation between two stations with the value of 0.732($p < 0.0001$), 0.753($p < 0.0001$), 0.845($p < 0.0001$) and 0.776($p < 0.0001$) respectively. The rest, Petaling Jaya-Kajang and Kajang-Shah Alam showed moderate correlation with the value of 0.678($p < 0.0001$) and 0.602($p < 0.0001$), respectively (Table 4). Klang (S1) was shown to be the main contributor to the PM₁₀ level and variation within 10 years period. Based on Malaysian Environmental Quality Report 2004 by Department of Environment, PM₁₀ were originating from stationary sources where industries as the highest contributor followed by power plants and the mobile sources from motor vehicles activity [42].

Table 3. Spearman correlation of PM₁₀ between Klang Valley regions for 10 years

Year	Variables	Klang	Petaling Jaya	Kajang	Shah Alam
2000-2009	Klang	1			
	Petaling Jaya	0.732	1		
	Kajang	0.753	0.678	1	
	Shah Alam	0.845	0.776	0.602	1

Values in bold are different from 0 with a significance level alpha = 0.05

Table 4. Probability value of the spearman correlation of PM₁₀ between Klang Valley regions for 10 years

Year	Variables	Klang	Petaling Jaya	Kajang	Shah Alam
2000-2009	Klang	0			
	Petaling Jaya	< 0.0001	0		
	Kajang	< 0.0001	< 0.0001	0	
	Shah Alam	< 0.0001	< 0.0001	< 0.0001	0

Values in bold are different from 0 with a significance level alpha = 0.05

Air quality classification using principal component analysis (PCA)

PCA has been applied for identifying the most and less significant location that contributes to air quality in Klang Valley region. Two types of test; Keiser-Meyer-Olkin Measure (KMO) and Barlett’s Test of Sphericity have been used in order to assess the sufficiency of the data [13, 37, 38]. KMO test shows the multicollinearity or in simple words, identification of similarity in correlation value exists between two or more items [15]. If there is a similarity in correlation between the items, those items will be measured in similar aspect. The benchmark’s value for KMO test is agreed to be more than 0.5 [37, 38]. The KMO value for this study is 0.726 which indicate that there is no serious multicollinearity problem for the Klang Valley’s PM₁₀ data and suitable to construct PCA. Besides, Barlett’s sphericity test also used in identification of the correlation between items and sufficiency of the data to construct the PCA with the benchmark value of less than 0.05 [13, 37, 38]. The result of Barlett’s test showed that the observed PM₁₀ data are sufficient to construct PCA with the obtained p-value less than 0.0001. These two sets of test are shown in Table 5.

Table 5. Keiser-Meyer-Olkin Measure (KMO) and Barlett’s Test of Sphericity

Types of PCA Test	Test’s Variable	Value
Keiser-Meyer-Olkin measure of sampling adequacy		0.726
Barlett’s Test of sphericity	Chi-square (observed values)	467.771
	Significant (p-value)	p < 0.0001

The Eigenvalue of these data is 3.391 and meet the PCA criterion because of the value larger than one (Eigenvalue > 1). The Eigenvalue shows the proportion of variance in every factor where the variance is significantly increase with increasing number of Eigenvalue [38]. Only 1 Eigenvalue was observed for the PCA analysis as shown in Table 6 because only PM₁₀ observation data were used in this study. The scree plot in Figure 3 can prove that only single Eigenvalue involve in this PCA interpretation and the factor that give a large different in variance will be shown clearly by the scree plot [38].

From this study, all of the air quality stations in Klang Valley affected by anthropogenic activity as seen in the F1. From Table 6, it shows that all of the stations gave the positive and strong factor loading [Klang (0.994), Petaling Jaya (0.902), Kajang (0.914) and Shah Alam (0.923)]. Therefore, over past 10 years period (2000 – 2009), particulate matter less than 10 micron (PM₁₀) is one of the most air pollutant that contributed to air quality problem in Klang Valley region. Strong positive factor loading lead to the source of particulate matter from traffic emission in which be the major source of PM₁₀ from automobiles’ fuel combustion activity [37, 43]. Figure 4 shows the dot matrix position base on factor loading of each air quality stations.




Figure 3. Scree plot of eigenvalue vs cumulative variability (%)

Table 6. Principal component analysis (PCA) interpreted result of PM₁₀ data in Klang Valley

Air Quality Station	Factor Loading			
	F1	F2	F3	F4
Klang	0.944	0.141	-0.234	0.185
Petaling Jaya	0.902	-0.246	0.347	0.076
Kajang	0.914	0.356	0.152	-0.123
Shah Alam	0.923	-0.256	-0.250	-0.142
Eigenvalue	3.391	0.273	0.261	0.076

The principal components were then being rotated to gain better understanding and interpretation of the data. Varimax rotation was used as a criterion [37] as proposed by other author [44], with only involve the consideration of component with an eigenvalue larger than 1. The variability and cumulative variability for rotated principal component scored both 84.773%. Table 7 shows the percentage of variability and cumulative variability of the PM₁₀ data after being rotated with Varimax rotation. On the other hand, Table 8 shows the percentage contribution of PM₁₀ in Klang, Petaling Jaya, Kajang and Shah Alam, which are 26.27%, 23.99%, 24.62% and 25.10% respectively. Therefore, it proved that the above statement can be accepted with the average percentage contribution of 25% for each area.

Table 7. Rotated percentage of variability and cumulative variability after Varimax rotation

	F1	F2	F3	F4
Variability (%)	84.773	6.818	6.516	1.893
Cumulative variability (%)	84.773	91.591	98.107	100.000




Figure 4. Dot matrix position base on factor loading of each air quality stations in Klang Valley region.

Table 8. Contribution of the variables after Varimax rotation

Contribution (%)	D1
Klang	26.27
Petaling Jaya	23.99
Kajang	24.62
Shah Alam	25.10
Average contribution	25.00
Total contribution	100.00

Air quality pattern using cluster Analysis (CA)

This study focused on four monitoring stations around Klang Valley region, where particulate matter less than 10 microns (PM₁₀) has been chosen for analysis. CA was used as an approach to find and put them into groups depending on their dissimilarity characteristics. PM₁₀ pattern which obtained from PCA output have been classified into three (3) different classes; Class 1, Class 2 and Class 3 based on location and can be illustrated in dendrogram such as in Figure 5.

Class 1 refers to air monitoring station in Sekolah Menengah Perempuan Raja Zarina, Klang (CA0011) representative as Klang air monitoring station. This station contributes highest number of dissimilarity to the graph where it is performed alone and giving significant indicator of air quality. In other words, Klang can be represented as a not good air condition. Class 2 covered two different air quality stations which are located in Sekolah Kebangsaan Seri Petaling, Petaling Jaya (CA0016) represent as Petaling Jaya air monitoring station and Country Height, Kajang (CA0023) represent Kajang air monitoring station. Class 3 refers to Sekolah Kebangsaan Taman Tun Dr. Ismail, Shah Alam (CA0025) which represents Shah Alam air monitoring station.

Klang (Class 1) is located near to industrial area and a busy port area originated at Port Klang, while the other two clusters are situated near to both industrial and residential area. The study showed that the sources of pollutant in all stations are affected by motor vehicles, industrial and factories activity. The level of PM₁₀ in Klang showed slightly higher concentration due to the port activity which was not found in the other stations [45].




Figure 5. Dendrogram view of cluster analysis for particulate matter (PM₁₀) in Klang Valley

Characteristic of PM₁₀ pattern within the Klang Valley region can be identified more using this CA method. This further analysis was discussed based on classes and the groups that might be produced. Figure 6 shows the pattern of PM₁₀ for three different clusters in Klang Valley based on year.

Throughout the 10 years period from 2000 to 2009, Class 1 (Klang) produces three patterns of group; Group 1 (2008 and 2009) and Group 2 (2002 and 2005). Meanwhile Group 3 (2000, 2001, 2003, 2004, 2006 and 2007) had shown the highest similarity within the year. Based on year 2000 to 2009, Class 2 which refers to Petaling Jaya and Kajang produces four different patterns of group; Group 1 (2008), Group 2 (2009), Group 3 (2003, 2004 and 2005) and lastly Group 4 (2000, 2001, 2002, 2006 and 2007). Class 3 (Shah Alam) produce 3 patterns, Group 1 (2000, 2001, 2006 and 2007), Group 2 (2008 and 2009) and Group 3 (2002, 2003, 2004 and 2005). The above finding shows that the concentration of PM₁₀ in the air were the highest at each station; Klang, Petaling Jaya, Kajang and Shah Alam at year 2008 and 2009.




Figure 6. Dendrogram (a – c) view of PM₁₀ pattern for three different clusters in Klang Valley based on year

Conclusion

Based on the result, significant relationship of particulate matter (PM₁₀) in Klang Valley, Malaysia and its pattern can be identified using statistical and envirometric technique. The relation between two variables can be determined by using Spearman Correlation Test. Strong correlations (> 0.7 with the p-value < 0.001) were obtained for all stations in determination of significant relationship between 2 stations except for Petaling Jaya-Kajang and Kajang-Shah Alam which recorded moderate correlation value 0.678(p < 0.001) and 0.602 (p < 0.001) respectively. Principal component analysis (PCA) was able to identify the most and less significant location which contributing to the level of PM₁₀. The study found that all four location have been affected by PM₁₀ which was determined as one of the air pollutant that contributes to the deterioration of air quality in Klang Valley. On other hand, Cluster analysis (CA) on the other hand was able to classify the PM₁₀ into three groups depending on their dissimilarities characteristic. PM₁₀ pattern which obtained from PCA output have been classified into three (3) different classes, Class 1 (Klang), Class 2 (Petaling Jaya and Kajang) and Class 3 (Shah Alam) based on location and possible period of extreme air quality degradation could be identified. These statistical and envirometric techniques have proved the impact of the various location on increased concentration of PM₁₀.

Acknowledgement

Gratitude and appreciation to Malaysian Department of Environment (DOE) for their provision of air quality data, and also to supervisor, committee members and colleagues for their efforts in assisting and completing this article.

References

1. Schwartz, J., Dockery, D. W., Neas, L. M., (1996). Is daily mortality associated specifically with fine particles. *Journal of the Air and Waste Management Association* 46(10): 927–939.
2. Shao, Y. P., (2000). Physics and modelling of wind erosion. Atmospheric and oceanographic sciences library, Sydney.
3. Abas, M. R. B., Oros, D. R. and Simoneit, B. R. T. (2004). Biomass burning as the main source of organic aerosol particulate matter in Malaysia during haze episodes. *Chemosphere*, 55: 1089 – 1095.
4. Afroz, R., Hassan, M. N. and Ibrahim, N. A. (2003). Review of air pollution and health impacts in Malaysia. *Environmental Research*. 92: 71 – 77.
5. Azmi, S. Z., Latif, M. T., Ismail, A. S., Juneng, L., Jemain, A. A. (2010). Trend and status of air quality at three different monitoring stations in the Klang Valley, Malaysia. *Air Quality, Atmosphere and Health*, 3: 53 – 64.
6. Gurjar, B. R., Butler, T. M., Lawrence, M. G. and Lelieveld, J. (2008). Evaluation of emissions and air quality in megacities. *Atmospheric Environment*, 42: 1593 – 1606.
7. Ozden, O., Dogeroglu, T. and Kara, S., (2008). Assessment of ambient air quality in Eskisehir, Turkey. *Environment International*, 34: 678 – 687.
8. Nawrot, T. S., Torfs, R., Fierens, F., De Henauw, S., Hoet, P. H. and Van Kersschaever, G. (2007). Stronger associations between daily mortality and fine particulate air pollution in summer than in winter: Evidence from a heavily polluted region in Western Europe. *Epidemiology and Community Health*, 61:146 – 149.
9. Mott, J. A., Mannino, D. M., Alverson, C. J., Kiyu, A., Hashim, J., Lee, Tee., Falter, K., Redd, S. C. (2005). Cardiorespiratory hospitalizations associated with smoke exposure during the 1997 Southeast Asia forest fires, *International journal of Hygiene and Environmental Health*, 2008: 75 – 85.
10. Baldasano, J. M., Valera, E., Jiménez, P. (2003). Air quality data from large cities. *The Science of The Total Environment*, 307: 141 – 165.
11. Azid, A., Juahir, H., Amran, M. A., Suhaili, Z., Osman, M. R., Muhamad, A., Abidin, I. Z., Sulaiman, N. H., Saudi, A. S. M. (2015). Spatial air quality modelling using chemometrics techniques: a case study in Peninsular Malaysia. *Malaysian Journal of Analytical Sciences*. 19(6): 1415 – 1430.
12. Azid, A., Juahir, H., Ezani, E., Toriman, M. E., Endut, A., Rahman, M. N. A., Yunus, K., Kamarudin, M. K. A., Hasnam, C. N. C., Saudi, A. S. M. and Umar, R. (2015b). Identification source of variation on regional impact of air quality pattern using chemometric. *Aerosol and Air Quality Research*, 15: 1545 –1558.
13. Azid, A., Juahir, H., Toriman, M. E., Endut, A., Kamarudin, M. K. A., Rahman, M. N. A., Hasnam, C. N. C., Saudi, A. S. M., Yunus, K. (2015). Source apportionment of air pollution: A case study in Malaysia. *Jurnal Teknologi*, 72(1): 83 – 88.

14. Amran, M. A., Azid, A., Juahir, H., Toriman, M. E., Mustafa, A. D., Hasnam, C. N. C., Azaman, F., Kamarudin, M. K. A., Saudi, A. S. M. and Yunus, K. (2015). Spatial analysis of the certain air pollutants using environmetric techniques. *Jurnal Teknologi*, 75(1): 241 – 249.
15. Isiyaka, H. A., Juahir, H., Toriman, M. E., Azid, A., Gasim, B. M. and Kamarudin, M. K. A. (2015). Assessment of the spatial variation and source apportionment of air pollution based on chemometric techniques: A case study in the Peninsular Malaysia. *Jurnal Teknologi*, 77(1): 33 – 44.
16. Isiyaka, H. A. and Azid, A. (2015). Air quality pattern assessment in Malaysia using multivariate techniques. *Malaysian Journal of Analytical Sciences*, 19(5): 966 – 978.
17. Azid, A., Juahir, H., Toriman, M. E., Kamarudin, M. K. A., Saudi, A.S.M., Hasnam, C. N. C., Aziz, N. A. A., Azaman, F., Latif, M. T., Zainuddin, S. F. M., Osman, M. R. and Yamin, M. (2014). Prediction of the level of air pollution using principal component analysis and artificial neural network techniques: A case study in Malaysia. *Water Air Soil Pollution*, 225(8): 1 – 14.
18. Bereton, R. (2003). Chemometrics. Data analysis for the laboratory and chemical plant. Wiley, Chichester.
19. Massart, D. L. and Kaufman, L. (1983). The interpretation of analytical chemical data by the use of cluster analysis, Wiley, New York.
20. Singh, K. P., Malik, A., Mohan, D. and Sintha, S., (2004). Multivariate statistical techniques for the evaluation of spatial and temporal variations in water quality of Chomti River (India) – a case study. *Water Research*, 38(18): 3980 – 3992.
21. Singh, K. P., Malik, A. and Sintha, S., (2005). Water quality assessment and apportionment of pollution sources of Gomti River (India) using multivariate statistical techniques – A case study. *Analytica Chimica Acta*, 538 (1-2): 355 – 374.
22. Kannel, P. R., Lee, S. R., Kanel, S. R. and Khan., S. P. (2007). Chemometrics application in classification and assessment of monitoring locations of an urban river system. *Analytica Chimica Acta*, 582: 390 – 399.
23. McKenna Jr., J. E. (2003). An enhance cluster analysis program with bootstrap significance testing for ecological community analysis. *Environmental Modelling and Software*, 8(2): 205 – 220.
24. Abas, M. R. and Simoneit, B. R. T. (1996). Composition extractable organic matter of air particle from Malaysia: initial study. *Atmospheric Environment*, 30(15): 2779 – 2793.
25. Omar, N. Y. M. J., Abas, M. R. B., Ketuly, K. A. and Tahir, N. M.(2002). Concentration of PAHs in atmospheric particle (PM₁₀) and road side soil particles collected in Kuala Lumpur, Malaysia. *Atmospheric Environment*, 36(2): 247 – 254.
26. Department of Environment (2009), Environmental quality report, Department of Environment Malaysia. Malaysia. Kuala Lumpur.
27. Tsikriktsis, N. (2005). A review of techniques for treating missing data in OM survey research. *Journal of Operations Management*, 24(1): 53 – 62.
28. Chan, L. S. and Dunn, O. J. (1972). The treatment of missing values in discriminant analysis. I. The sampling experiment. *Journal of the American Statistical Association*, 67: 473 – 477.
29. Chan, L. S., Gilman, J. A. and Dunn, O. J. (1976). Alternative approaches to missing values in discriminant analysis. *Journal of the American Statistical Association*, 71: 842 – 844.
30. Raymond, M. R. and Roberts, D. M., (1987). A comparison of methods for treating incomplete data in selection research. *Educational and Psychological Measurement*, 47: 13 – 26.
31. Saudi, A. S. M., Juahir, H., Azid, A., Toriman, M. E., Kamarudin, M. K. A., Saudi, M. M., Mustafa, A. D. and Amran, M. A. (2015). Flood risk pattern recognition by using environmetric technique: A case study in Langat river basin. *Jurnal Teknologi*, 77(1): 145–152.
32. Saudi, A. S. M., Azid, A., Juahir, H., Toriman, M. E., Amran, M. A., Mustafa, A. D., Azaman, F., Kamarudin, M. K. A. and Saudi, M. M. (2015). Flood risk pattern recognition using integrated chemometric method and artificial neural network: A case study in the Johor river basin. *Jurnal Teknologi*, 74(1): 165 – 170.
33. Saudi, A. S. M., Juahir, H., Azid, A., Kamarudin, M. K. A., Kasim, M. F., Toriman, M. E., Aziz, N. A. A., Hasnam, C. N. C. and Samsudin, M. S. (2015). Flood risk pattern recognition using chemometric technique: A case study in Kuantan river basin. *Jurnal Teknologi*, 72(1): 137 – 141.
34. Chua, Y. P. (2013), Spearman correlation test; Book 3, basic research statistic, analysis of likert scale data, Mc Graw-Hill Education (Ed. 2): pp. 242 – 262.

35. Kamaruddin, A. F., Toriman, M. E., Juahir, H., Zain, S. M., Rahman, M. N. A., Kamarudin, M. K. A., and Azid, A. (2015). Spatial characterization and identification sources of pollution using multivariate analysis at Terengganu river basin, Malaysia. *Jurnal Teknologi*, 77(1): 269 – 273.
36. Mutalib, S. N. S. A., Juahir, H., Azid, A., Sharif, S. M., Latif, M. T., Aris, A. Z., Zain, S. M. and Dominick, D. (2013). Spatial and temporal air quality pattern recognition using environmetric techniques: A case study in Malaysia. *Environmental Science: Processes Impacts*, 15: 1717 – 1728.
37. Ul-Saufie, A. Z., Yahya, A. S., Ramli, N. A., Rosaida, N., and Hamid, H. A., (2013). Future daily PM₁₀ concentration prediction by combining regression models and feedwork backpropagation models with principal component analysis (PCA). *Atmospheric Environment*, 77: 621 – 637.
38. Chua, Y. P. (2009), Factor analysis and cluster analysis; Book 5, regression test, factor analysis and SEM analysis, Mc Graw-Hill Education: pp. 170 – 299
39. Abdul-Wahab, S. A., Bakheit, C.S. and Al-Alawi, S. M. (2005). Principal component and multiple regression analysis in modelling of ground-level ozone and factors affecting its concentrations. *Environmental Modelling Software*, 20: 1263 – 1271.
40. Markou, M. and Kassomenos, P. (2010). Cluster analysis of five years of back trajectories arriving in Athens, Greece. *Atmospheric Research*, 98: 438 – 457.
41. Masiol, M., Rampazzo, G., Ceccato, D., Squizzato, s., and Pavoni, B. (2010). Characterization of PM₁₀ sources in coastal area near Venice (Italy): An application of factor-cluster analysis. *Chemosphere*, 80: 771 – 778.
42. Department of Environment (2011). Environmental Quality Report, Department of Environment Malaysia. Malaysia. Kuala Lumpur.
43. Ghazali, N. A., Ramli, N. A. and Yahya, A. S. (2009). A study to investigate and model the transformation of nitrogen dioxide into ozone using time series plot. *European Journal of Scientific Research* 37(2): 192 – 205.
44. Kaiser, H. F. (1958). The varimax criterion for analytic rotation in factor analysis. *Psychometrika*, 23: 187 – 200.
45. Dominick, D., Juahir, H., Latif, M. T., M. Zain, S. and Aris, A. Z. (2012). Spatial assessment of air quality patterns in Malaysia using multivariate analysis. *Atmospheric Environment*, 60: 172 – 181.