

26TH MEDICAL & HEALTH RESEARCH WEEK

**“COLLABORATIVE FRONTIERS IN MEDICAL RESEARCH:
UNITING MINDS FOR SCIENTIFIC BREAKTHROUGHS”**

Program & Abstract

5-9 AUGUST 2024

**LECTURE HALL 2, LEVEL 2, PRECLINICAL BLOCK
FACULTY OF MEDICINE
UNIVERSITI KEBANGSAAN MALAYSIA**

ORGANISER

SPI

Sekretariat
Penyelidikan
& Inovasi

Secretariat of Research & Innovation

TABLE OF CONTENT

Message from the Dean of Faculty of Medicine	2
Message from the Deputy Dean (Research & Innovation)	3
Organizing Committee	4
List of Speakers	6
List of Facilitators	7
Tentative Program	8
List of Abstracts	11
Abstracts	14

**MESSAGE
DEAN FACULTY OF MEDICINE
UNIVERSITI KEBANGSAAN MALAYSIA**



Assalamualaikum warahmatullahi wabarakatuh

It is a great honour to welcome you to the 26th Annual Medical and Health Research Week. I am delighted to see such an exceptional scientific gathering of minds dedicated to advancing the frontiers of medical and health research. This event promises an outstanding value of knowledge and fosters collaboration, all aimed at achieving research excellence and advancing global healthcare.

In the first place, I want to acknowledge Prof. Dr. Mohammad Nasir Shafiee's leadership for the enduring continuation of Medical and Health Research Week over the years. My heartfelt gratitude is also dedicated to Dr Muhammad Luqman Nasaruddin, the chairman of this year, his organising committee, and the Secretariat of Research & Innovation, whose hard work and thorough planning have made this event happen. Well done, all.

Recently, UKM secured an impressive position in the global higher education landscape, achieving a world ranking of 136th. This outstanding accomplishment determines our dedication, excellence, and innovation. Securing a position in the top 150 universities worldwide is no small feat. It reflects our commitment to delivering high-quality education, fostering groundbreaking research, and contributing significantly to our community and beyond.

As such, our Faculty of Medicine is among the highest contributors to the UKM MyRA point; considering our research outputs and their impact on global knowledge are pivotal in UKM's world ranking. Our commitment to addressing pressing global challenges through innovative research contributes to the betterment of society and the advancement of medical sciences.

We are proud of the achievements of our faculty members in research and innovation throughout the years. A number of our lecturers have recently been recognised as being among the World's Top 2% of Scientists. This recognition enhances the reputation of our institution on the global stage.

In addition to this impressive accomplishment, it is important to highlight another significant achievement of our faculty members. Their remarkable ability to secure grants from international agencies, such as the Natural Science Foundation of Guangdong, China and the Health Research Board of Ireland. Not to forget, industry grants from Merck Sharp & Dohme United States, Bayer AG Germany and AstraZeneca United Kingdom. This achievement speaks volumes about their expertise, the quality of their research, and their standing within the global scientific community. This level of support is crucial for conducting high-impact research that can lead to groundbreaking discoveries and significant advancements.

In today's rapidly evolving field of medicine, it has become increasingly evident that no single medical discipline is capable of providing all the answers or solutions to the complex and multifaceted health challenges we face. Each medical specialty brings its own unique expertise and perspective to the table, highlighting the importance of transdisciplinary collaboration. Through collaboration, combining diverse viewpoints, sharing knowledge, and uniting our efforts, we can make cutting-edge findings and tackle the complex challenges in the healthcare system. This is the impactful power of collaboration, consistent with this year's theme.

I encourage all participants to take this unique opportunity to actively engage in the workshops and immerse in insightful discussions during Research Week. Interact with experienced scholars during the networking sessions and gain insights from their achievements. Your enthusiasm, dedication, and fresh perspectives are essential to driving innovation and shaping the future of medical research.

Let's begin this event with an open heart and a mutual commitment to furthering medical and health research. Thank you for your presence and participation, and I look forward to a fruitful week.

Prof. Datin Dr. Marina Mat Baki
Dean Faculty of Medicine
Universiti Kebangsaan Malaysia

**MESSAGE
DEPUTY DEAN (RESEARCH & INNOVATION)
FACULTY OF MEDICINE
UNIVERSITI KEBANGSAAN MALAYSIA**



Assalamualaikum Warahmatullahi Wabarakatuh and Salam Sejahtera,

A warm welcome to each of you for the 26th Annual Medical and Health Research Week. It is with great enthusiasm and pleasure that I stand before you as the Advisor of this event. This week promises to be a beacon of collaboration and innovation, full of inspiration towards research excellence.

This year's theme, "Collaborative Frontiers in Medical Research: Uniting Minds for Scientific Breakthrough," highlights the power of teamwork in driving innovation and discovery in research. The fact that you are here today is evidence of how seriously you take your commitment, despite your hectic schedule, to pushing the boundaries of knowledge and making a positive impact in medical research.

Throughout this week, we have meticulously curated a diverse program of activities, including talks, interactive workshops, and panel discussions. Let's not forget the e-posters showcasing research findings, with some of the studies showing collaboration internationally. Congratulation! Each session is designed to encourage intellectual exchange, inspire creativity, and build a robust collaboration. We hope you make the most of this event by networking across academic boundaries.

This event is more than just a congregation of brilliant minds; it is a testament to strengthening collaboration and innovation to advance research. This week is a valuable opportunity for our postgraduate students, lecturers, and young scientists for their research growth, mentorship, and networking. Connect with experienced researchers, engage in stimulating discussions, and draw lessons from their successes. Take this opportunity to ignite your passion, expand your horizons, and advance research excellence.

Finally, I want to acknowledge the incredible efforts of our organising committee members under the leadership of Dr. Muhammad Luqman Nasaruddin, who have worked tirelessly to make this event possible. Their dedication, creativity, and meticulous planning have paved the way for an extraordinary week.

Ladies and gentlemen, thank you for being here. Let's begin this journey by bringing together minds toward scientific advancements. Make this week a memorable celebration of knowledge, progress, and the endless possibilities in medical science. By working together, we can establish connections, foster synergy, and discover the endless potential of our collective experience. Thank you.

Prof. Dr. Mohammad Nasir Shafiee
Deputy Dean (Research & Innovation)
Faculty of Medicine
Universiti Kebangsaan Malaysia

ORGANIZING COMMITTEE

ADVISOR

Prof. Dr. Mohammad Nasir Shafiee
(Deputy Dean of Research and Innovation, Faculty of Medicine)

CHAIRMAN

Dr. Muhammad Luqman Nasaruddin

DEPUTY CHAIRMAN

Assoc. Prof. Dr. Aneeza Khairiyah Wan Hamizan

SECRETARY

Mr. Muhammad Khaidir Hashim

TREASURER

Mrs. Nurazilah Mohamed Sallehuddin

ASISTANT TREASURER

Mrs. Aznida Hasan

PROTOCOL COMMITTEE

Dr. Muhammad Luqman Nasaruddin
Dr. Maziah Ahmad Marzuki
Dr. Isma Liza Mohd Isa
Mr. Muhammad Khaidir Hashim

PROMOTION COMMITTEE

Dr. Daniel Law Jia Xian
Mr. Razif Adli Rosli
Mrs. Ruzanna Nizar

PROGRAM COMMITTEE

Dr. Wong Sok Kuan
Dr. Farah Hanan Fathihah Jaffar
Mrs. Siti Noorain Hamid
Mr. Nik Mohd Nashrullah Nik Mohd Fadzil

SMALL GROUP DISCUSSION COMMITTEE

Assoc. Prof. Dr. Mohd.Heikal Mohd Yunus
Dr. Ekram Alias
Mr. Nik Mohd Nashrullah Nik Mohd Fadzil

POSTER COMMITTEE

Assoc. Prof. Dr. Wong Yin Ping
Dr. Tan Shing Cheng
Dr. Nurul 'Izzah Ibrahim
Dr. Wong Kon Ken
Dr. Rizal Abd Rani
Dr. Nur Fatin Nabilah Mohd Sahardi
Ms. Nazratul Fareha Shari

REGISTRATION COMMITTEE

Mrs. Nurazilah Mohamed Sallehuddin
Ms. Nur Izyani Abd Razak

SOUVENIRS COMMITTEE

Mrs. Ruzanna Nizar
Mrs. Aznida Hasan

TECHNICAL COMMITTEE

Mr. Abdul Malek Mohd Yunan
Mr. Nik Mohd Nashrullah Nik Mohd Fadzil
Mr. Muhammad Khaidir Hashim

LIST OF SPEAKERS

Speaker	Title
Assoc. Prof. Dr. Jayakumar Murthy	A Step-by-Step Guide to Writing Your Literature Review
Prof. Emeritus Dato' Dr. Wan Zurinah Wan Ngah	Developing & Submitting a Project Proposal
Assoc. Prof. Dr. Wan Haslina Wan Abdul Halim	Clinical Trial and Randomization
Assoc. Prof. Dr. Chin Kok Yong	Writing a Scoping Review: Tips and Challenges
Assoc. Prof. Dr. Isa Naina Mohamed Prof. Dr. Mohd Shahrir Mohamed Said	Avoiding & Controlling for Biases Ethics in Clinical Research
Assoc. Prof. Dr. Azimatun Noor Aizuddin	- Study Designs: Action Research, Case Study, Causal, Cohort, Cross-Sectional, Descriptive - Study Designs: Experimental, Exploratory, Historical, Longitudinal, Observational, Philosophical, Sequential
Mrs Qurratul Syaheera Ahmad Termizi Dr. Zuraidah Che' Man	Managing your References - Sample Size Calculation - Exploratory Data Analysis
Assoc. Prof. Dr. Norfazilah Ahmad	- Testing Hypothesis - Chi Square & Other Qualitative Data Analysis
Prof. Dr. Azmawati Mohammed Nawi	T-test & ANOVA
Assoc. Prof. Dr. Nazarudin Safian	- Correlation & Regression - Non-parametric Tests - Agreement Analysis & Kappa

LIST OF FACILITATORS

Assoc. Prof. Dr. Azimatun Noor Aizuddin
Assoc. Prof. Dr. Chin Kok Yong
Assoc. Prof. Dr. Jaya Kumar A/L Murthy
Assoc. Prof. Dr. Mohd Faizal Ahmad
Assoc. Prof. Dr. Norliza Muhammad
Assoc. Prof. Dr. Yogeswaran Lokanathan
Assoc. Prof. Dr. Daniel Law Jia Xian
Assoc. Prof. Dr. Noor Akmal Shareela Ismail
Assoc. Prof. Dr. Hanani Abdul Manan
Assoc. Prof. Dr. Aneeza Khairiyah Wan Hamizan
Assoc. Prof. Dr. Nor Rafeah Tumian
Dr. Nor Haslinda Abd Aziz
Dr. Ekram Alias
Dr. Fazlina Nordin
Dr. Nurul 'Izzah Ibrahim
Dr. Rizuana Iqbal Hussain
Dr. Wong Sok Kuan
Dr. Vinoth a/l Kumarasamy
Dr. Nadiah Sulaiman

TENTATIVE PROGRAM

Module A

5/8/2024 (Monday)	Program	Venue
Moderator: Assoc. Prof. Aneeza Khairiyah Wan Hamizan / Dr. Nurul 'Izzah Ibrahim		
8:00 am – 8:15 am	Registration and Attendance	Lecture Hall 2, Preclinical Building
8:15 am – 9:15 am	A Step-by-Step Guide to Writing Your Literature Review by Assoc. Prof. Dr. Jayakumar Murthy	
9:15 am – 10:15 am	Developing & Submitting a Project Proposal by Prof. Emeritus Dato' Dr. Wan Zurinah Wan Ngah	
10:15 am – 10:30 am	Break	Preclinical lobby, Level 1
10:30 am – 11:30 am	Clinical Trial and Randomization by Assoc. Prof. Dr. Wan Haslina Wan Abdul Halim	Lecture Hall 2, Preclinical Building
11.30 am – 1.30 pm	Writing a Scoping Review: Tips and Challenges by Assoc. Prof. Dr. Chin Kok Yong	
1:30 pm – 2:30 pm	Break and Lunch	
2:30 pm – 4:30 pm	Developing Your Proposal: Hands on Session with Facilitators (Day 1)	BPK Level 5, Preclinical Building

Module B & Opening Ceremony

6/8/2024 (Tuesday)	Program	Venue
Moderator: Dr. Farah Hanan Fathihah Jaffar		
8:15 am – 8:30 am	Registration and Attendance	Lecture Hall 2, Preclinical Building
8:30 am – 9:30 am	Avoiding & Controlling for Biases by Assoc. Prof. Dr. Isa Naina Mohamed	
9:30 am – 10:30 am	Ethics in Clinical Research by Prof. Dr. Mohd Shahrir Mohamed Said	
10:30 am – 11:00 am	Break and VIP arrival	Preclinical lobby, Level 1
11:00 am – 11.05 am	Opening Ceremony Singing of National Anthem & Doa Recitation	Lecture Hall 2, Preclinical Building
11.05 am – 11.10 am	Speech by the Deputy Dean (Research & Innovation) by Prof. Dr. Mohammad Nasir Shafiee	
11.10 am – 11.30 am	Speech by the Dean of Faculty of Medicine by Prof. Dr. Marina Mat Baki	
11:30 am – 11:35 am	Montage of Research Week	
11.35 am – 11.45 am	Introduction to Researcher of the Year by Prof. Dr. Raja Affendi Raja Ali	
11:45 am – 11:50 am	Video of Researcher of The Year	
11:50 am – 12:00 pm	Award giving and Photo session	
12.00 pm – 1.00 pm	Talk by Researcher of the Year by Prof. Dr. Norfilza Mohd Mokhtar	
1.00 pm – 2:30 pm	Break and Lunch	Preclinical lobby, Level 1
2:30 pm – 4:30 pm	Developing Your Proposal: Hands on Session with Facilitators (Day 2)	BPK Level 5, Preclinical Building

Module C

7/8/2024 (Wednesday)	Program	Venue
Moderator: Dr. Ekram Alias		
8:00 am – 8:15 am	Registration and Attendance	Lecture Hall 2, Preclinical Building
8:15 am – 9:15 am	Study Designs: Action Research, Case Study, Causal, Cohort, Cross-Sectional, Descriptive by Assoc. Prof. Dr. Azimatun Noor Aizuddin	
9:15 am – 10:15 am	Study Designs: Experimental, Exploratory, Historical, Longitudinal, Observational, Philosophical, Sequential by Assoc. Prof. Dr. Azimatun Noor Aizuddin	
10:15 am – 10:30 am	Break	Preclinical lobby, Level 1
10:30 am – 12:00 pm	Poster Presentation	Student Lounge, Level 2 Preclinical Building
10:30 am – 11:30 am	Managing your References by Puan Quratul Syaheera Ahmad Termizi	Lecture Hall 2, Preclinical Building
11:30 am – 12:30 pm	Sample Size Calculation by Dr Zuraidah Che' Man	
12:30 pm – 1:30 pm	Exploratory Data Analysis by Dr Zuraidah Che' Man	
1:30 pm – 2:30 pm	Break and Lunch	Preclinical lobby, Level 1
2:30 pm – 4:30 pm	Developing Your Proposal: Hands on Session with Facilitators (Day 3)	BPK Level 5, Preclinical Building

Module D

8/8/2024 (Thursday)	Program	Venue
Moderator: Assoc. Prof. Dr. Law Jia Xian		
8:15 am – 8:30 am	Registration and Attendance	Lecture Hall 2, Preclinical Building
8:30 am – 9:15 am	Testing Hypothesis by Assoc. Prof. Dr. Norfazilah Ahmad	
9:15 am – 10:00 am	Chi Square & Other Qualitative Data Analysis by Assoc. Prof. Dr. Norfazilah Ahmad	
10:00 am – 10:15 am	Break	Preclinical lobby, Level 1
10:15 am – 11:00 am	T-test & ANOVA by Prof. Dr. Azmawati Mohammed Nawi	Lecture Hall 2, Preclinical Building
11:00 am – 11:45 am	Correlation & Regression by Assoc. Prof. Dr. Nazarudin Safian	
11:45 am -12:30 pm	Non-Parametric Tests by Assoc. Prof. Dr. Nazarudin Safian	
12:30 pm – 1:15 pm	Agreement Analysis & Kappa by Assoc. Prof. Dr. Nazarudin Safian	
1:15 pm – 2:30 pm	Break and Lunch	Preclinical lobby, Level 1
2:30 pm – 4:30 pm	Developing Your Proposal: Hands on Session with Facilitators (Day 4)	BPK Level 5, Preclinical Building

Module E

9/8/2024 (Friday)	Program	Venue
8:45 am – 9:00 am	Registration and Attendance	Lecture Hall 2, Preclinical Building
9:00 am – 12:00 pm	Proposal Presentation	
	Moderator: Assoc. Prof. Dr. Mohd Heikal Yunus Group 1 – 9 Lecture Hall 1, Preclinical Building	Moderator: Dr. Ekram Alias Group 10 – 19 Lecture Hall 2, Preclinical Building
12:00 pm – 12:30 pm	Prize Giving and Closing	Lecture Hall 2, Preclinical Building
12:30 pm – 2:45 pm	Lunch	Preclinical lobby, Level 1

LIST OF ABSTRACT

No	Code	Title	Authors	Page No
CLINICAL				
1	C1	A "Starter Pack" for Big Data Analytics in Preventive Medicine – A Systematic Review	Sheng Qian Y ¹ , Nur Alya Asyiqin IS ¹ , Hui Ying C ¹ , Azizah Z ¹ , Zhi Qin O ¹	14
2	C2	Prevalence Of Treatment Failure and its Associated Factors Among Syphilis Patients In Hospital Canselor Tuanku Muhriz: A Retrospective Study	Afifah Aqilah AM ^{1,4} , Najma K ^{1,4} , Asrul AW ^{2,4} , Mohd Rizal AM ^{3,4} , Petrick P ^{1,4}	15
3	C3	Prevalence and Effectiveness of Health Promotion Programs for Preventing Opisthorchis viverrini Infection in Nakhon Phanom Province, Thailand	<u>Khanittha Nakharangsu</u> ¹ , Zulkarnain Md Idris ² , Sriwipa Chuangchaiya ^{1*}	16
4	C4	A Pilot Study on Stature Estimation Using The Sternum in a Malaysian Population	<u>Tay Cheong W</u> ¹ , Mohamed Swarhib S ¹ , Faridah MN ¹	17
5	C5	Expression of PDL-1 by Immunohistochemistry in mature T cell & T/NK cell lymphoma and its correlation with the prognostic outcomes	Nabil Aljunid ¹ , Nurwahyuna Rosli ¹ , Nor Rafeah Tumian ² , Azimatun Noor ³ , Nur Maya Sabrina Tizen Laim ¹ , Muatamarulain Mustangin ¹ , Suria Hayati Md Pauzi ¹	18
6	C6	The Dual Dilemma: Hypercalcaemia and Persistent PTH Elevation After Parathyroidectomy – Unmasking FHH as a Second Pathology	Jia Wen S ¹ , Wan Muhammad Azfar WS ¹ , Noor Ayuni MP ² , Ana Daliela M ²	19
7	C7	Entomotoxicological Analysis Using Third Instar Larvae in a Postmortem Case	<u>Hafiz WM</u> ^{1,2*} , Izzul M ³ , Faridah MN ¹ , Heo CC ⁴	20
8	C8	Perception Of Teachers On Adaptive Skills Among Preschoolers: A Qualitative Study	Masne K ^{1,2*} , Ainna ZM ¹ , Nor ARR ^{1,2} , Farahiyah WY ^{1,2}	21
9	C9	Effectiveness of Active Release Technique on Posture and Neck Range of Motion in College Students with Upper Cross Syndrome	Mgt Mohd Hafiz, MH ¹ , Azira Iqlima, R ¹ , Nurhanisah, S ¹ , Mohd Murtadha, R ¹ , Santibuana, AR ^{1*}	22
10	C10	Awareness of and Attitudes Toward Pharmacogenetic Testing for HLA-B*15:02 among Parents/Guardians of Epileptic Patients in Malaysia: A Qualitative Study	<u>Iman Zaki</u> ¹ , Adli Ali ^{1*} , Sharifah Azween Omar ¹ , Wong Sau Wei ¹ , Nor Asyikin Tahir ² , Shamin Saffian ² , A'dina Rizal ³ , Nurazwa Rizal ³	23
11	C11	Trans-anal Barotrauma in Children: A Rare Consequence of Playful Insufflation	Tan Chun Chau ¹ , Munirah Osman ² , Aireen Zamhot ³ , <u>Evelyn Chau Yi Wen</u> ^{3*}	24
12	C12	Diagonal Earlobe Crease: Preliminary Study for Its Significance in Cardiovascular-related Deaths	<u>Bashirah Mohd Nor</u> ^{1*} , Wan Mohammad Hafiz Wan Razali ^{1,2} , Tay Cheong Woo ¹ , Faridah Mohd Nor ¹ , Mohamed Swarhib Shafee ¹	25
13	C13	Screen Time In Children With Autism Spectrum Disorder Of School-Going Age	<u>Lolyasmin Lokman</u> ¹ , <u>Norazlin Kamal Nor</u> ^{2*}	26
14	C14	Solid Pseudopapillary Neoplasm: A Rare Presentation	<u>Vannese Phey Jia Wong</u> ¹ , Jievashini V. Murali ¹ , Guang Hong Ong ¹	27
15	C15	The Venomous Sight Killer	<u>YeeLin Lee</u> ¹ , Alex Chau Sim Yee ² , LayHui Ng ³ , Siti Zakiah Binti Md Khair ⁴ , Muhammad Aiman Bin Mohd Suhaimi ⁵	28

16	C16	Health related quality of life and recovery following cardiac surgery via median sternotomy	Md Ali Nur Ayub ^{1*} , Abdul Rashid Muhammad Fathi ² , Abdul Rahman Mohd Ramzisham ¹ , Ismail Muhammad Ihsamuddin ¹ , Abdul Manaf Mohd Rizal ³ , Md Ali Katijahbe ⁴	29
17	C17	Hand Grip Strength among Year 1 and Year 2 Undergraduate Students of Faculty of Medicine and Health Sciences, Universiti Malaysia Sarawak (UNIMAS)	Saut Alexander Sureng ¹ , Syamsul-Amri Aida Nurlateefah ¹ , Puang Ivy Juliette Yi Hui ¹ , Azizi Marsha Amirah ¹ , Fong Isabel Lim ² , Sahdi Haniza ^{3*}	30
18	C18	Prediction of Treatment Response in Brain Metastasis Patients treated with Gamma Knife Using Radiomics and Machine Learning from Pre-Treatment MRI Scans	Abdulrahman U ^{1,*} , Noorazrul Y ² , Hanani AM ³ , Ramesh KAK ⁴ , Siti Khadijah H ⁴	31
19	C19	Prediction of local recurrence in head and neck cancer following chemotherapy using radiomic features from pre-radiotherapy MRI	Wun Chin Leong ^{1,4} , Hanani Abdul Manan ² , Caryn Chan Mei Hsien ³ , Yoke Fui Wong ⁴ , Eriananiza Mohd Salim ⁵ , Noorazrul Yahya ¹	32
20	C20	Let's get personal! Molecular testing for targeted therapies in solid tumours: HCTM experience	Nurwahyuna R ^{1*} , Nur Fa'izah AM ² , Noor Azlin MH ³	33
21	C21	Necrotizing Soft Tissue Infection: Microbiological Distribution From District Region	Amri Dahdi ¹ , Yazid Bajuri ² , Norhaslinda Bahaudin ³	34
22	C22	Pre-Radiotherapy Balloon Dilatation of the Eustachian Tube: Reducing Radiation-Induced Dysfunction in Nasopharyngeal Carcinoma Patients	Lorna Ting KN ^{1,2} , Noor Dina H ¹ , Lum SG ¹ , Siti Hajar S ² , Mohd Razif MY ¹ , Khairiyah S ³	35
23	C23	Perception Of Parents On Adaptive Skills Among Preschoolers: A Qualitative Study	Masne K ^{1,2*} , Nur Subhanie M A ¹ , Nor Affi R ^{1,2} , Farahiyah WY ^{1,2}	36
24	P1	Understanding Children's Play Perceptions and Play Preferences in Sarawak: A Qualitative Study	Erza Nurfarziany Zalani ^{1*} , Farahiyah Wan Yunus ¹	37
25	P2	A Qualitative Study of the Effect of Dialysis on the Performance of Daily Living Activities in Diabetic Clients in the Kuala Selangor Region	Atikah Suib ¹ , Farahiyah Wan Yunus ¹	38
26	P3	The Quality of Life and Sleep Quality of Health Sciences Undergraduate Students	Noor Azliyana Azmi ¹ , Farahiyah Wan Yunus ¹	39
27	P4	Practice and Perceived Challenges among Clinical Immunologists Toward Genetic Testing and Genetic Counseling in Malaysia: A Qualitative Study	Vigneswary Balakrishnan ¹ , Sharifah Azween Syed Omar ¹ , Nadiah Sulaiman ² , Chai-Eng Tan ³ and Adli Ali ^{1*}	40
28	P5	Reliability, Usability, and Feasibility of the MINAT App among Children Aged 6 to 12 in Malaysia	Wan Sarah Hanan Wan Mohd Hilmi ¹ , Nor Affi Razaob ^{1,2} , Masne Kadar ^{1,2} , Farahiyah Wan Yunus ^{1,2}	41
29	P6	Effects of Different Footwear Types and Barefoot on Static and Dynamic Balance among College Students	Nurfarah Najihah, KN ¹ , Azira Iqlima, R ¹ , Nurhanisah, S ¹ , Mohd Murtadha, R ¹ , Santibuana, AR ^{1*}	42
30	P7	Validity, Reliability and Profile of Children Handwriting Based on OTulis in Malaysia	Nurnabihah Hashim ¹ , Nor Affi Razaob ^{1,2} , Masne Kadar ^{1,2} , Farahiyah Wan Yunus ^{1,2}	43
31	P8	Validity and Profile of Children's Activity Preferences of 'MINAT'	Wei Shan Tan ^{1*} , Nor Affi Razaob ^{1,2} , Masne Kadar ^{1,2} , Farahiyah Wan Yunus ^{1,2}	44
32	P9	MicroRNA Expression Pattern in Complete and Partial Hydatidiform Moles	Wai Kit Chia ^{1,2} , Hannah Xin Yi Yeoh ¹ , Wei-Kang Lee ³ , Kok Jun Liew ³ , Chee Choong Hoh ³ , Teck Yee Khong ^{4,5} , Yin Ping Wong ^{1*} , Geok Chin Tan ^{1*}	45

33	P10	The Effects of Calcium Phosphate Cement Doped with Palm Tocotrienol on Tibial Bone Defect in Ovariectomised Rats through Static Bone Histomorphometry	<u>Siti Sarah MD</u> ¹ , Wong SK ^{1*} , Chin KY ¹ , Fairus A ²	46
34	P11	Relationship Between Level of Self Satisfaction and Self Confident Among Medical Student at National University of Malaysia (UKM) Towards Simulation Based Learning	<u>W. Nurfaeana</u> , Mohammad Arif Kamarudin, Alias Mahmud*	47
35	P12	Exploring the Dose-Dependent Impact of Celastrol on Anthropometric Data and Blood Glucose Levels in ApoE-knockout Mice Fed a High-Fat Diet	<u>Arifah, AD</u> ¹ , Thuhairah Hasrah, AR ¹ , B. Vimala, RMTB ² , Nasibah, A ^{1*}	48
36	P13	Optimising Caco-2 Cells and Bifidobacterium Bifidum Co-Culture to Investigate the Possible Mechanism of Lapatinib-Induced Diarrhoea	<u>Nur Syafina, H</u> ¹ , Wan Nor Izzah, WMZ ^{1*} , Nur Ain Najiha, R ¹ , Jesmine, K ¹ , Narimah, AHH ¹ and Hassanain Issam, YT ²	49
37	P14	Ultrastructural characteristics of placentas in pregnancies complicated with fetal growth restriction	<u>Che En Tan</u> ¹ , Kevin Boon Ping Ku ¹ , Wirda Indah Farouk ² , Rahana Abd Rahman ³ , Geok Chin Tan ^{1*} , Yin Ping Wong ^{1*}	50
38	P16	Mapping Regorafenib's Protein Targets in Colorectal Cancer: Insights from Molecular Fingerprinting	<u>Farina Syazwani Md Azhan</u> , Muhammad Redha Abdullah Zawawi, Teck Yew Low, Pey Yee Lee*	51
39	P17	Enrichment of GBP7 Mutation in Patients with Primary Aldosteronism	<u>Fatin Athirah Pauzi</u> ¹ , Nurfatin Syazreen Mod Rizam ¹ , Munkit Choy ² , Simon Williams ² , Muaatamarulain Mustangin ³ , Geok Chin Tan ³ , Xilin Wu ⁴ , Elisabeth Ng ^{4,5} , Kate Laycock ⁴ , Azraai Bahari Nasruddin ⁶ , Norlela Sukor ^{1,7} , Morris Jonathan Brown ⁴ , Bernard Keaveney ² , Elena Aisha Azizan ^{1*}	52
40	P18	Exploring NPHS Mutations in Pediatric Patients with Congenital and Steroid-Resistant Nephrotic Syndrome	<u>Jun Xin Lee</u> , Noor Akmal Shareela binti Ismail*	53
41	P19	Effects of Kelulut Honey on the Frontal Lobe and Cerebellum in Amyloid β -induced Alzheimer's Disease Rat Model	<u>Suhilen TS</u> ¹ , Shaik A ¹ , Teoh SL ¹ , Yahaya MF ^{1*}	54
42	P20	Proteomic profiling reveals interferon pathway upregulation in FOLFOX-resistant colorectal cancer cells	<u>Yeoh Yeelon</u> , Low Teck Yew, Nadiah Abu, Lee Pey Yee*	55
43	P21	Precision Medicine Unveils the Molecular Profiles of Extracellular Matrix Anabolism, Inflammation, and Nociception Tailored to the Severity of Degenerative Disc Disease in Humans	<u>Nurul Fariha Z</u> ^{1,2} , Izzat Z ² , Kieran J ^{2,3} , Karina D ⁴ , Aoibhinn K ⁴ , Ahmad Hafiz K ⁵ , Muhamad Nazrin M ⁵ , Sabarul Afian M ⁵ , Elvy Suhana M R ¹ , Fairus A ¹ , David P F ^{2,4} , Abhay P ² , Aiden D ³ , Isma Liza M I ^{1,2,4}	56

C1

A “Starter Pack” for Big Data Analytics in Preventive Medicine – A Systematic Review

Sheng Qian Y, Nur Alya Asyiqin IS, Hui Ying C, Azizah Z, Zhi Qin O*Department of Public Health Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latiff, Bandar Tun Razak, 56000 Cheras, Kuala Lumpur, Malaysia*

ABSTRACT

Big data has transformative potential in preventive medicine. However, clinicians often experience difficulties when performing big data analytics due to the requirement of multidisciplinary skill sets and vast knowledge on information technology. To overcome these challenges, the current review aimed to summarise the various data sources, algorithms, software, and statistical analyses used in big data analytics in disease prevention. Literature search was performed using PubMed, Scopus, Web of Science, and Institute of Electrical and Electronics Engineers databases. Studies were included if they included big data analytics, reported on primary, secondary or tertiary prevention of disease, published 2010-2024, as well as written in English language. Among the 31 included studies, authors had utilised data from electronic medical records (n = 11), nationwide census/registries (n = 7), and insurance databases (n = 4). Machine learning algorithms were mostly used, including random forest (n = 11), logistic regression (n = 10), and gradient boosting (n = 7). Big data analyses were mostly conducted using R software (n = 15), Python (n = 6), and SAS (n = 6). Area under the receiver operating characteristic curve (n = 11), sensitivity/recall (n = 10), positive predictive values (PPV)/precision (n = 8), specificity (n = 6), and F1-score (n = 6) were the commonly reported model outputs. In summary, clinicians who plan to conduct big data analytics shall have access to publicly available datasets, obtain basic knowledge on machine learning algorithms as well as to master the operating and interpretative skills using various big data software.

Keywords: Big data analytics; machine learning; preventive medicine

C2

Prevalence of Treatment Failure and its Associated Factors among Syphilis Patients in Hospital Canselor Tuanku Muhriz: A Retrospective Study

Affah Aqilah AM^{1,4}, Najma.K^{1,4}, Asrul AW^{2,4}, Mohd Rizal AM^{3,4} Petrick P^{1,4}

¹Department of Medicine, ²Department of Medical Microbiology and Immunology, ³Department of Community Health, ⁴ Faculty of Medicine, Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia

*Corresponding: fifamalik@gmail.com

ABSTRACT

The increasing prevalence of syphilis and its limited information in Malaysia is a public health concern. The objectives of this study is to determine the prevalence of treatment failure and its potential associated factors among syphilis patients in Hospital Canselor Tuanku Muhriz (HCTM), the clinical stages of syphilis during the first presentation in our hospital and if there was any difference in clinical presentation between human immunodeficiency virus (HIV) and non-HIV patients with syphilis. The study was conducted over 5 years period in HCTM in patients who were serologically confirmed to have syphilis by positive treponemal antibody test. Their data were retrieved to obtain information on socio-demographics, comorbidities, clinical characteristics, treatment received, and treatment outcome using a convenient sampling method. Logistic regression analysis was performed to determine the independent factors associated with treatment failure in syphilis patients. Total of 209 patients who were confirmed to have syphilis were screened but only 141 patients were analysed. Most of the syphilis patients were young adults, male, HIV positive, men-sex-men (MSM), asymptomatic and presented as latent syphilis. There was no difference in the clinical presentation of syphilis between HIV and non-HIV patients. 119 patients (84.4%) experienced treatment success, whereas 22 patients (15.6%) experienced treatment failure. Among patients with syphilis, females ($p < 0.001$), HIV status ($p = 0.018$), and asymptomatic patients on routine screening ($p = 0.020$) were the initial predictors of treatment failure. Females were 4.3 times more likely to have treatment failure [95%CI: 1.08,16.8]; $p = 0.039$) as compared with male patients. In conclusion, female patients should be monitored closely for treatment failure after syphilis treatment. Further larger multicenter studies should be done to better describe those who have a higher risk of treatment failure and whether there are any long-term clinical implications.

Keywords: Syphilis; HIV; prevalence; treatment failure' retrospective study

C3

Prevalence and Effectiveness of Health Promotion Programs for Preventing *Opisthorchis viverrini* Infection in Nakhon Phanom Province, ThailandKhaniittha Nakharangsu¹, Zulkarnain Md Idris², Sriwipa Chuangchaiya^{1*}¹Department of Community Health, Faculty of Public Health, Kasetsart University, Chalermphrakiat Sakon Nakhon Province Campus, Sakon Nakhon, Thailand²Department of Parasitology and Medical Entomology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

*Correspondence: sriwipa.c@ku.th

ABSTRACT

Opisthorchis viverrini is one of the endemic helminths in Thailand. Although several public health initiatives have been implemented to reduce opisthorchiasis, the prevalence remains high in endemic areas. This study aimed to determine the prevalence of *O. viverrini* infection and evaluate the effectiveness of health promotion programs in opisthorchiasis prevention among rural communities in Northeast Thailand. A cross-sectional study was conducted between May and December 2019 in Tha Uthen district in Nakhon Phanom Province, Thailand. Stool samples were collected from 310 individuals (≥ 18 years) and processed using the Kato-Katz technique to determine the presence of *O. viverrini* and other intestinal helminths. Health literacy programs were created to understand the knowledge, attitude, and practice (KAP) of *O. viverrini* among the experimental (30 participants) and control (30 participants) groups. Participants' KAP levels were assessed using questionnaires before and after the intervention program (12 weeks). The prevalence of intestinal helminths and *O. viverrini* was 17% and 11%, respectively. Individuals aged between 40-59 years (adjusted odds ratio [aOR] = 0.38, $p = 0.038$) and those with a previous history of opisthorchiasis (aOR = 1.97, $p = 0.033$) were positively associated with a higher risk of *O. viverrini* infection. Regarding the health promotion programs, the experimental group showed a significant increase in average scores in KAP and preventive measures of *O. viverrini* compared to the control group ($p < 0.05$). This finding suggests that although opisthorchiasis is still prevalent in Nakhon Phanom, health promotion programs were effective in improving communities' KAP regarding the infection.

Keywords: Opisthorchiasis; epidemiology; prevention; risk factor; Thailand

C4

A Pilot Study on Stature Estimation Using the Sternum in a Malaysian Population

Tay Cheong W, Mohamed Swarhib S, Faridah MN

Forensic Unit, Department of Pathology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur, Malaysia.

*Correspondence: tcwoo137@gmail.com

ABSTRACT

Identification of deceased individuals is a critical aspect of examining unidentified human remains. Estimating stature aids forensic anthropologists in narrowing down potential matches among victims. This study aims to highlight the relationship between different measurements of the sternum and stature. It seeks to develop a specific linear regression formula for stature estimation in the Malaysian adult population using sternal lengths. This pilot study was conducted at the Forensic Unit, HCTM, where measurements of sternum and stature were collected from 15 males and 15 females during medicolegal autopsy. Initial findings revealed significant differences in various sternal lengths between males and females ($p < 0.05$). Median total sternal length (TSL) was notably longer in males, 205.400mm (IQR 30.540) compared to females, 177.950mm (IQR 18.890) ($p = 0.003$). Also, there were significant correlation between different sternal lengths and stature ($p < 0.001$). The highest observed correlation coefficient (r) was 0.79, which between stature, the length of the manubrium and sternal body (LMB), and TSL. Linear regression analysis demonstrated a significant linear relationship between sternal lengths and stature ($p < 0.001$). The highest observed regression coefficient of determination (R^2) was 0.61 when using TSL. This preliminary investigation suggests that various measurements of the sternum reliably predict stature and could be employed for stature estimation in the Malaysian adult population when long bones are not available for biological profiling of human skeletal remains. Further research with a larger sample size is necessary to confirm these findings and explore the potential use of sternal lengths for stature estimation in the identification of deceased individuals.

Keywords: Stature estimation; sternal lengths; Malaysian; anthropology; forensic.

C5

Expression of PDL-1 by Immunohistochemistry in Mature T cell & T/NK Cell Lymphoma and its Correlation with the Prognostic Outcomes

Nabil ALJUNID¹, Nurwahyuna ROSLI¹, Nor Rafeah TUMIAN², Azimatun NOOR³, Nur Maya Sabrina TIZEN LAIM¹, Muaatamarulain MUSTANGIN¹, Suria Hayati MD PAUZI¹

¹Department of Pathology, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, 56000, Kuala Lumpur, Malaysia

²Department of Internal Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, 56000, Kuala Lumpur, Malaysia

³Department of Public Health, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, 56000, Kuala Lumpur, Malaysia

ABSTRACT

Introduction: Programmed cell death 1 (PD-1)/programmed cell death ligand 1 (PD-L1) is a cell surface protein involved in normal immune checkpoint function. By binding to PD-1, PD-L1 provides a suppressive signal to T-cells and an anti-apoptotic signal to tumour cells, leading to T-cell dysfunction and tumour survival.

Methodology: PDL-1 expression was evaluated in various types of T/NK cell NHLs using PDL-1 antibody clone 22C3 on formalin-fixed paraffin-embedded tissue sections. Tumour proportion score (TPS) of 1% or more is taken as positive.

Results: 71.4% of 42 cases of T-cell NHLs were PDL-1 positive. There is no significant association between PDL1 expression with age, gender, race, LDH level, positive B symptoms, Ann Arbor stage, ECOG score, extranodal involvement and IPI score. Nevertheless, Ann Arbor stage, ECOG score and IPI risk groups showed a significant impact towards the survival outcomes. PDL1-positive patients have demonstrated a poorer overall survival (OS) and progression-free survival (PFS), however, were not statistically significant.

Discussion: PDL-1 positive T cell lymphoma showed a worse survival outcomes, suggesting PDL-1 positivity is an adverse prognostic factor. The lack of significant survival impact suggests that PDL-1 alone may not be a robust prognostic marker in T-cell lymphomas. Further studies with larger cohorts are necessary to clarify the role of PD-L1 in these malignancies and to improve the prognostic and therapeutic strategies for patients with T cell NHLs.

Keywords: PDL-1; PD-1; T/NK cell lymphoma; Non-Hodgkin lymphoma

C6

The Dual Dilemma: Hypercalcaemia and Persistent PTH Elevation after Parathyroidectomy – Unmasking FHH as a Second Pathology

Jia Wen S¹, Wan Muhammad Azfar WS¹, Noor Ayuni MP², Ana Daliela M²

¹Department of Pathology, Faculty of Medicine, Universiti Kebangsaan Malaysia

²Department of Pathology, Hospital Melaka

*Correspondence: p125808@siswa.ukm.edu.my

ABSTRACT

Introduction: Primary hyperparathyroidism (PHPT) is one of the main causes of hypercalcaemia. In contrast, familial hypocalciuric hypercalcaemia (FHH) is a less common diagnosis for patients with asymptomatic hypercalcaemia who usually have a positive family history. *Case report:* We present a case of a 53-year-old woman who presented with persistently elevated PTH levels and hypercalcaemia. She was diagnosed with PHPT and subsequently underwent parathyroidectomy. However, her serum calcium levels fluctuated between the upper half of the reference range and the range of hypercalcaemia post-surgery. Her PTH levels were persistently elevated despite normal total 25-hydroxyvitamin D levels, with overlapping calcium/creatinine clearance ratio (CCCR). Ultrasound of the neck and parathyroid gland was repeated twice post-surgery and showed no other parathyroid lesions. An alternative diagnosis of concurrent FHH was considered. *Discussion:* CCCR is a screening test for both disorders, albeit with limited sensitivity and specificity and a prerequisite of a good 24-hour urine collection. In our case, the inconsistent CCCRs did not help to indicate recurrent PHPT or newly diagnosed FHH. Further investigations with screening of family members and a mutation of the calcium-sensing receptor (CaSR) are necessary. *Conclusion:* Dual pathology should be considered if the patient does not respond to conventional treatment. After screening the family members, mutation analysis of the CaSR gene will be an important test to differentiate between PHPT and FHH.

Keywords: Hypercalcaemia; hyperparathyroidism; CCC; CaSR gene

C7

Entomotoxicological Analysis using Third Instar Larvae in a Postmortem CaseHafiz WM^{1,2*}, Izzul M³, Faridah MN¹, Heo CC⁴

¹*Forensic Unit, Department of Pathology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Cheras, 56000 Kuala Lumpur, Malaysia*

²*Department of Forensic Pathology, Faculty of Medicine, Sungai Buloh Campus, Universiti Teknologi MARA, 47000 Sungai Buloh, Selangor, Malaysia*

³*National Institute of Forensic Medicine Malaysia, 50586 Kuala Lumpur, Malaysia*

⁴*Institute of Medical Molecular Biotechnology, Faculty of Medicine, Sungai Buloh Campus, Universiti Teknologi MARA, 47000 Sungai Buloh, Selangor, Malaysia*

*Corresponding author: p103741@siswa.ukm.edu.my

ABSTRACT

Entomotoxicology is a branch of forensic entomology that studies the effects of drugs and toxins on insects that feed on decomposing human tissues. This field plays a crucial role in forensic investigations, especially in cases where traditional toxicological methods are limited or unavailable. This study aims to report a case of chemical content detection using third instar larvae collected from corpses. The maggot samples were collected from a decomposed body during a postmortem examination. The sample was sent to the Institute of Medical Molecular Biotechnology, Universiti Teknologi MARA Sungai Buloh, for entomotoxicology analysis using the third instar larvae analyzed with the Gas Chromatography-Mass Spectrometry to rule out drug-related death. The analysis showed that some suspicious compounds were detected in the insect's evidence samples and not in the control samples. Based on the Scientific Working Group for the Analysis of Seized Drugs, the compounds found were classified as ketone functional group, ganglionic blocker, opioid analgesic, antihistamine, histamine antagonist, and neuroprotective agent. The findings of a variety of analytes in a toxicological analysis, including compounds with different functional groups and pharmacological effects, can be indicative of the use of designer drugs, potential polydrug use, and the complexity of drug adulteration. In conclusion, advanced analytical techniques are essential for accurately identifying these substances and detection of these compounds can have significant legal and criminal implications, potentially indicating illicit drug use or aiding in criminal investigations where poisoning or drug-related foul play is suspected.

Keywords: Entomotoxicology; forensic entomology; Gas Chromatography-Mass Spectrometry; designer drugs; toxicological analysis

C8

Perception of Teachers on Adaptive Skills among Preschoolers: A Qualitative Study

Masne K^{1,2*}, Ainna ZM¹, Nor ARR^{1,2}, Farahiyah WY^{1,2}

¹Occupational Therapy Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia

²Centre for Rehabilitation & Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia

*Correspondence: masne_kadar@ukm.edu.my

ABSTRACT

Adaptive skills encompass social, self-care, and academic abilities crucial for children's daily functioning and overall development. While extensive research exists on adaptive skills in atypical children, studies on typical children in Malaysia are limited. Research exploring the link between teachers' perceptions and preschoolers' adaptive skills is scarce. Understanding these relationships is vital for coherent child development. This study aims to explore teachers' perceptions of adaptive skills in preschoolers. A phenomenological approach recruited six female teachers from the Klang Valley using purposive sampling. Inclusion criteria included a minimum of 3 years teaching experience, minimum of diploma qualification, and teaching typical children aged 3-6 years. Interviews were conducted online, and face-to-face, and thematic analysis was employed, ensuring validity through COREQ checklist criteria. The study identified three main themes regarding adaptive skills: exploring foundational aspects, key influences and strategies for nurturing them, as well as challenges faced in the process. It involved three private and three government preschools adhering to Ministry of Education standards. The participants highlighted the significance of self-care, socialization, and communication skills in child development, school readiness, and academic achievement. Challenges included inadequate facilities and diverse parent-teacher approaches. Effective teaching methods included interactive and play-based approaches. The study also noted that teachers often prioritized social and communication skills over holistic adaptive skill development. These findings underscore the importance of adaptive skill development from teachers' perspectives, addressing concepts, challenges, and experiences, that will guide practitioners in effectively supporting children's developmental milestones.

Keywords: Teacher's perception; adaptive skills; preschool children; communication skills; activities of daily living; social skill

C9

Effectiveness of Active Release Technique on Posture and Neck Range of Motion in College Students with Upper Cross Syndrome

Mgt Mohd Hafiz, MH, Azira Iqlima, R, Nurhanisah, S, Mohd Murtadha, R, Santibuana, AR*

Faculty of Pharmacy and Health Sciences, Universiti Kuala Lumpur, Royal College of Medicine Perak, Ipoh, Perak, Malaysia

*Correspondence: santibuana@unikl.edu.my

ABSTRACT

Upper cross syndrome (UCS) is a common postural dysfunction characterized by the imbalance of muscles around the neck and shoulder girdle. UCS is becoming increasingly prevalent among college students due to inactivity and prolonged periods of poor posture. However, studies on the physiotherapeutic strategies focusing on active release technique (ART) and ice pack therapy (IPT) for treating UCS especially among college students is still lacking. This study aimed to compare the effectiveness of ART and IPT on posture and neck range of motion (ROM) in college students with UCS. This experimental study involving 40 college students (mean age: 23.25 ± 1.68) using convenience sampling technique. Participants were randomly assigned into two groups: Group A as an intervention group received ART and Group B served as control group received IPT. Postural analysis and goniometer were used to assess posture and neck ROM. The data were analysed using paired t-test and independent t-test. There was a significant difference on posture and neck ROM for all variables in intervention group pre- and post-intervention of ART ($p < 0.05$). However, in control group, there was no significant difference on posture and some variables of neck ROM pre- and post-intervention of IPT ($p > 0.05$). The intervention group receiving ART showed statistically significant improvements in both posture and neck ROM compared to the control group receiving IPT ($p < 0.05$). This study proves that ART is more effective than IPT, suggesting that ART should be considered a preferred physiotherapeutic strategy for treating college students with UCS.

Keywords: Active Release Technique (ART); Ice Pack Therapy (IPT); posture analysis; Neck Range of Motion (ROM); Upper Cross Syndrome (UCS)

C10

Awareness of and Attitudes toward Pharmacogenetic Testing for HLA-B*15:02 among Parents/Guardians of Epileptic Patients in Malaysia: A Qualitative Study

Iman Zaki¹, Adli Ali^{1*}, Sharifah Azween Omar¹, Wong Sau Wei¹, Nor Asyikin Tahir², Shamin Saffian², A'dina Rizal³, Nurazwa Rizal³

¹Department of Paediatrics, Faculty of Medicine, Universiti Kebangsaan Malaysia

²Department of Pharmacy, Faculty of Pharmacy, Universiti Kebangsaan Malaysia

³Department of Biological Sciences and Biotechnology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia

*Correspondence: adli.ali@ppukm.ukm.edu.my

ABSTRACT

Pharmacogenetic testing for the HLA-B*15:02 allele is essential to prevent severe adverse reactions to carbamazepine in epileptic patients. Current research predominantly focuses on healthcare professionals' awareness and knowledge of pharmacogenetic testing, rather than the general public. The extent of knowledge among parents/guardians in Malaysia regarding their awareness and perceptions of HLA-B*15:02 pharmacogenetic testing remains unclear. This study explores the awareness, knowledge, attitudes, experiences, and gaps among parents/guardians of epileptic patients concerning HLA-B*15:02 pharmacogenetic testing. Qualitative methods were used, conducting semi-structured interviews with parents/guardians of children tested at the Neurology Clinics of UKM Specialist Children's Hospital. Of 30 potential participants, 11 were unreachable and 9 were unaware of the test, resulting in a cohort of 10 participants. Interviews were held via Zoom or face-to-face, and thematic analysis revealed 5 key themes and 15 sub-themes. The findings show generally low awareness and understanding of HLA-B*15:02 pharmacogenetic testing, often misunderstood as a blood test rather than a genetic test. Healthcare providers often did not provide specific genetic details, highlighting significant comprehension gaps. Participants expressed a need for more information on the testing procedure, potential outcomes, and implications. This study emphasizes the crucial role of comprehensive pre-test counseling in enhancing understanding and supporting informed decision-making among parents/guardians of epilepsy patients. Particularly noteworthy is the experience of two participants whose children experienced adverse drug reactions despite receiving negative test results. These insights aim to improve communication and education about pharmacogenetic testing, ultimately enhancing patient care and safety in clinical practice.

Keywords: Pharmacogenetics; HLA-B*15:02; epileptic patients; carbamazepine; adverse drug reactions

C11

Trans-anal Barotrauma in Children: A Rare Consequence of Playful InsufflationTan Chun Chau¹, Munirah Osman², Aireen Zamhot³, Evelyn Chau Yi Wen^{3*}¹*Department of Emergency Medicine, Hospital Seberang Jaya, Jalan Tun Hussein Onn, Seberang Jaya, Perai, Pulau Pinang, Malaysia*²*Department of Emergency Medicine, Hospital Sultanah Bahiyah Km 6, Jalan Langgar, Alor Setar, Kedah, Malaysia*³*Department of Emergency Medicine, Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Kuala Lumpur, Malaysia*Correspondence: evelynchau@ukm.edu.my**ABSTRACT**

Trans-anal barotrauma resulting from high-pressure air insufflation through anal orifice is extremely rare and typically associated with prank behaviour among young male industrial workers. In children, fewer than ten cases have been reported in the past decade. We report a case of a 10-year-old boy who presented with abdominal pain, distension, and scrotal swelling after his granduncle playfully introduced a tire air inflator into his anal orifice through his pants. Upon assessment in the Emergency Department, his vital signs were stable. Physical examination revealed widespread subcutaneous emphysema, extending from lower chest wall to bilateral scrotum. Blood investigations were normal and his erect chest x-ray confirmed pneumoperitoneum. Anorectal examination under general anaesthesia showed erythema with superficial lacerations. Sigmoidoscopy showed intact mucosa without tears or ecchymosis up to 35cm from the anal verge. Despite these injuries, the child remained hemodynamically stable and was treated non-surgically. Given the unusual nature of injury, the Suspected Child Abuse and Neglect Team was involved for comprehensive evaluation and intervention. He was later discharged well to his mother. Trans-anal barotrauma from high-pressured air can cause a spectrum of colorectal injuries, ranging from minor serosal tears to full-thickness perforations and intra-abdominal compartment syndrome. Severity depends on air pressure, flow velocity, anal resting pressure and distance from the source to anus. Obtaining detailed history from both the child and family is crucial for prompt diagnosis and intervention. Education on proper use of mechanical equipment is also essential, as most incidents are unintentional and result from lack of awareness.

Keywords: Barotrauma; child neglect; insufflation; pneumoperitoneum; trans-anal

C12

Diagonal Earlobe Crease: Preliminary Study for Its Significance in Cardiovascular-related Deaths

Bashirah Mohd Nor^{1*}, Wan Mohammad Hafiz Wan Razali^{1,2}, Tay Cheong Woo¹,
Faridah Mohd Nor¹, Mohamed Swarhib Shafee¹

¹*Department of Pathology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, Kuala Lumpur, Malaysia*

²*Department of Forensic Pathology, Faculty of Medicine, Sungai Buloh Campus, Universiti Teknologi MARA, Jalan Hospital, Sungai Buloh, Selangor, Malaysia*

*Correspondence: p109433@siswa.ukm.edu.my

ABSTRACT

Diagonal earlobe crease (DELC), also known as Frank's sign, is a skin manifestation associated with coronary artery disease. Previous autopsy studies have established a connection between DELC and cardiovascular (CV) deaths. This pilot study in Malaysia aimed to determine the prevalence of DELC in autopsy cases of CV deaths compared to non-CV deaths. A cross-sectional study was conducted with 10 autopsy cases of CV deaths and 10 cases of non-CV deaths at the Forensic Unit, Hospital Canselor Tuanku Muhriz. The presence of DELC was evaluated using ear photographs and categorized into mild, moderate, or severe grades for both ears. The study population had a median age of 46.5 years (IQR 28.0), with 85% males and 15% females. The ethnic distribution was 35% Malay, 50% Chinese, and 15% Indian. A significant difference in DELC prevalence was found between CV and non-CV deaths ($p=0.033$). Also, a significant difference in DELC prevalence was observed between cases with critical coronary artery stenosis and those without ($p=0.008$). CV deaths exhibited higher median DELC grades in both ears, with p -values of 0.05 and 0.023 for the right and left ears, respectively. These results suggest that DELC is a valuable marker for coronary artery stenosis and may aid in predicting CV deaths.

Keywords: Earlobe crease; cardiovascular death; coronary artery disease; Frank's sign

C13

Screen Time in Children with Autism Spectrum Disorder of School-Going AgeLolyasmin Lokman¹, Norazlin Kamal Nor^{2*}¹Department of Paediatric, Faculty of Medicine, Universiti Kebangsaan Malaysia²Hospital Pakar Kanak-kanak, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur*Correspondence: norazlyn@ppukm.ukm.edu.my**ABSTRACT**

Autism spectrum disorder (ASD) is a neurodevelopmental disorder characterized by deficits in social communication and interaction and restrictive and repetitive behaviours, interest or activities. Children with ASD have been reported to be more interested in electronic forms of entertainment and communication compared to human interaction. In this cross-sectional study, we assessed the screentime habits in ASD children as well as the association between excessive screentime duration and sleep, physical activity, behaviour and obesity in school-aged children with ASD. We recruited 91 ASD children aged 6-12 years old attending Child Development Centre, Hospital Pakar Kanak-Kanak (HPKK) Universiti Kebangsaan Malaysia (UKM) over 16 months. Parents completed demographic questionnaire as well as validated Children's Sleep Habits Questionnaire (CHSQ) assessing sleep disturbance, Physical Activity Questionnaire (PAQ-C) assessing physical activity and ASD-Behaviour Problem for Children questionnaire (ASD-BPC), assessing challenging behaviours. BMI of the children were obtained. Chi-square testing was performed to determine association between excessive screentime and sleep disturbance, adequacy of physical activity and obesity. The mean duration of screentime per day was 2.96 hours (sd 1.61), with 28.6% having screentime of more than 4 hours per day ("excessive screentime"). Predictors of excessive screen time among our ASD patients included low paternal education level, personal ownership of gadgets, and lack of parental control. Longer screen time ($r = 0.225$, p -value = 0.015) and shorter sleeping hours ($r = -0.22$, p -value = 0.036) were correlated with more severe behavioural problems. In addition, higher levels of physical activity was associated with reduced sleep disturbances (p -value = 0.048). Understanding this may help paediatricians give suitable advice to parents in managing their children with ASD holistically.

Keywords: Autism; screen time; sleep habit; physical activity; behaviour

C14

Solid Pseudopapillary Neoplasm: A Rare Presentation

Vannese Phey Jia Wong, Jievashini V. Murali, Guang Hong Ong

Hepato-Pancreato-Biliary Unit, Department of General Surgery, Hospital Sultanah Bahiyah, Alor Setar, Kedah

**Correspondence: vannese.pjwong@gmail.com*

ABSTRACT

Solid pseudopapillary neoplasm (SPN) is a rare tumour which accounts for about 1-3% of exocrine tumour. It has low malignant potential and high survival rate which occur in young women. We present a case of 18-year-old who presented with epigastric mass for one year associated with constitutional symptoms and abdominal pain. CT abdomen reported as large pancreatic cyst causing portal vein compression and patient underwent percutaneous drainage in which the content was haemorrhagic fluid. Ca 19-9 was 5.5 and fluid for amylase showed 806 and the cytology result is negative. The size of cystic mass appeared similar in the repeated CT and subsequently patient underwent open distal pancreateo-splenectomy and noted huge tumour measuring 15 x 15 cm at distal pancreas. Histopathology revealed features of solid pseudopapillary neoplasm. Patient responded well to the treatment and did not required chemotherapy. The overall prognosis of SPN is good however it is difficult to diagnose it as patient normally presented with non-specific symptoms. It is a case was initially thought to be pseudocyst, but the drainage fluid show differently. SPN of pancreas can be confused as pseudocysts hence young female who presented with any big pancreatic lesion should be considered as SPN.

Keywords: Solid pseudopapillary neoplasm; pancreatic pseudocyst

C15

The Venomous Sight Killer

YeeLin Lee, Alex Chau Sim Yee, LayHui Ng, Siti Zakiah Binti Md Khair, Muhammad Aiman Bin Mohd Suhaimi

Hospital Pakar Sultanah Fatimah, Jalan Salleh, Taman Utama Satu, 84000, Muar, Johor

**Correspondence: yeelin_525@hotmail.com*

ABSTRACT

Bee sting toxic keratopathy is an uncommon ocular injury caused by the injection of venom to cornea. It can result in various complications with triad of penetrating, immunologic and toxic effects of the stinger and its injected venom. This is an ocular emergency. This is a case report of a 65 years old malay man, underlying dyslipidemia presented with history of trauma to his right eye from a bee. He presented 2 days after the trauma with complain of severe right eye pain and redness associated with reduced vision. His presenting visual acuity was right eye counting finger, left eye 6/9 without relative afferent pupillary defect. Anterior segment of the right eye showed injected conjunctiva with generalized corneal edema and descemet folds. There was a retained stinger embedded intrastromally located at periphery 6 o'clock with surrounding infiltrate. Anterior chamber appeared deep, unable to visualize cells due to hazy cornea, no hypopyon. Intraocular pressure was 22mmHg. Fundus details unable to visualize however B-scan showed retina was flat. Left eye examination was normal. He was started with intensive topical steroids, topical antibiotics and cycloplegics over his right eye. The retained stinger was removed surgically under local anesthesia together with corneal suturing. Post-operatively, his right eye vision improved to 6/18 (pinhole 6/9) with corneal scarring. His medications were tapered down gradually over a two-month period. Bee sting toxic keratopathy are of rare occurrence, but are associated with potential severe visual impairment. Early recognition and prompt treatment are crucial to prevent significant ocular morbidity.

Keywords: Bee sting; toxic keratopathy

C16

Health related quality of life and recovery following cardiac surgery via median sternotomy

Md Ali Nur Ayub^{1*}, Abdul Rashid Muhammad Fathi², Abdul Rahman Mohd Ramzisham¹, Ismail Muhammad Ihsamuddin¹, Abdul Manaf Mohd Rizal³, Md Ali Katijjahbe⁴

¹*Cardiothoracic Unit, Department of Cardiothoracic Surgery, Hospital Canselor Tuanku Muhriz*

²*Department of General Surgery, Hospital Canselor Tuanku Mukhriz*

³*Department of Community Health, Hospital Canselor Tuanku Muhriz*

⁴*Department of Physiotherapy, Hospital Canselor Tuanku Muhriz*

*Correspondence: drfathirashid@gmail.com

ABSTRACT

Recovery and health-related quality of life (HRQOL) is becoming increasingly recognized for determining surgical success from the perspectives of patients and healthcare providers. The aim of this study is to assess changes in HRQOL up to 6 months after cardiac surgery performed using a median sternotomy. This is a prospective cohort observational study. A total of 38 participants following median sternotomy completed the assessment of HRQOL using EQ-ED-5L at baseline (pre-operatively), prior to discharge, at 4-6 weeks, 3 and 6 months post-operatively. There was significant improvement in HRQOL after median sternotomy with an effect size of 0.38 ($P < 0.001$). The mean (SD) for the EQ-5L-5D was 61.9 (SD 21.7) at baseline; prior to discharge was 54.3 (SD 19.4), 65.8 (SD 24.5) at four weeks and 72.1 (SD 27.4) at three months and 75.4 (SD 28.3) 6 months respectively. All patients showed substantial improvement thereafter at the third month, crossed the baseline and steadily increased at the sixth month. Cardiac surgery via median sternotomy is associated with significant improvement as assessed via EQ-ED-5L. The majority of participants subjectively perceived low on HRQOL but tended to improve over time. This is the first prospective study on patient-reported outcomes in cardiac surgery patients in Malaysia. Our results may assist clinicians to interpret changes in HRQOL using EQ-5D-5L across time after median sternotomy. This result can be used to justify the impact of surgery, promotes shared decision-making, and facilitates patient-centered care at both clinical and policy levels, especially in light of an ageing population in Malaysia

Keywords: Health related quality of life; sternotomy; cardiac surgery

C17

Hand Grip Strength among Year 1 and Year 2 Undergraduate Students of Faculty of Medicine and Health Sciences, Universiti Malaysia Sarawak (UNIMAS)

Saut Alexander Sureng¹, Syamsul-Amri Aida Nurlateefah¹, Puang Ivy Juliette Yi Hui¹, Azizi Marsha Amirah¹, Fong Isabel Lim², Sahdi Haniza^{3*}

¹Medical student, Faculty of Medicine and Health Sciences, Universiti Malaysia Sarawak (UNIMAS)

²Department of Paraclinical Sciences, Faculty of Medicine and Health Sciences, Universiti Malaysia Sarawak (UNIMAS)

³Department of Orthopaedics, Faculty of Medicine and Health Sciences, Universiti Malaysia Sarawak (UNIMAS)

*Correspondence: shaniza@unimas.my

Hand grip strength (HGS) is crucial as stronger HGS in healthcare trainees is linked to better performance of clinical skills requiring manual dexterity. However, there is limited research on HGS determinants among medical and nursing students in Malaysia. This study aims to investigate the associations between HGS and gender, little finger flexor digitorum superficialis (FDS) muscle functionality, body mass index (BMI), physical activity frequency, hand dominance, and ethnicity of preclinical students enrolled in the Doctor of Medicine (MD) and Nursing programmes at Faculty of Medicine and Health Science (FMHS), Universiti Malaysia Sarawak (UNIMAS). Data were gathered via Google forms for socio-demographic information, anthropometric measurements for BMI calculation, the Short International Physical Activity Questionnaire (IPAQ) for physical activity levels, and the Modified Baker's test for little finger FDS functionality. HGS was measured using a Camry electronic hand dynamometer following American Society of Hand Therapists (ASHT) guidelines. Statistical Package for Social Sciences (SPSS) version 29.0 was utilised for data analysis, employing descriptive statistics and inferential tests with significance set at $p < 0.05$. This study found that HGS was significantly higher in males, greater BMI groups, and higher physical activity levels. Right-handed individuals demonstrated significantly stronger HGS in their dominant hands, whereas left-handed individuals displayed more balanced strength between hands. Ethnicity and presence or absence of the little finger FDS had no significant effect on HGS. Overall, this study offers insights into the factors influencing HGS, highlighting the importance of gender, hand dominance, physical activity levels, and BMI.

Keywords: Hand grip strength (HGS) little finger flexor digitorum superficialis (FDS); hand dominance Body Mass Index (BMI)

C18

Prediction of Treatment Response in Brain Metastasis Patients treated with Gamma Knife Using Radiomics and Machine Learning from Pre-Treatment MRI Scans

Abdulrahman U^{1,2*}, Noorazrul Y², Hanani AM³, Ramesh KAK⁴, Siti Khadijah H⁴

¹Department of medical radiography, Ahmadu Bello University Zaria, Nigeria

²Diagnostic Imaging and Radiotherapy, CODTIS, Faculty of Health Sciences, National University of Malaysia, Jalan Raja Muda Aziz 50300 Kuala Lumpur, Malaysia

³Department of Radiology, Hospital Pakar Kanak Kanak (UKM Specialist children's hospital) Universiti kebangsaan Malaysia, Jalan yaacob Latif, Bandar Tun Razak, Kuala Lumpur Malaysia

⁴Gamma Knife Centre, Faculty of Medicine Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

*Corresponding author: abdulumaru08@gmail.com, P130570@siswa.ukm.edu.my

ABSTRACT

Predicting treatment outcome early after Gamma Knife radiosurgery (GKRS) is important and may improve patient outcome. Radiomics, which utilizes a quantitative approach in image analysis, has the potential to provide more powerful and accurate predictors for treatment response when combined with ML. This study aims to utilize MRI radiomics and ML to predict treatment response of brain metastases (BM) following GKRS. The study was approved by the ethical committee of UKM (UKM.FSK.PNI.800-2/27/9(JEP-2022-524)). Tumors were classified as good response when more than 25% reduction in size. Radiomics features were extracted from 108 tumors in 30 patients that meet the inclusion criteria from April 2019 to March 2024. The ROIs were segmented manually using the planning system, and the final outline was approved by an experienced neurosurgeon. The images were resampled to 1x1x1 mm³ as recommended by Image Biomarker Standardization Initiative (IBSI), and radiomic features were extracted using 64 bin size with mean relative intensity ± 3 sd using LifeX software (v7.6.0). K-nearest neighbours (KNN) model was used with Synthetic Minority Oversampling Technique (SMOTE) to address data imbalance. A total of 16 features were selected using a 2-step feature selection (correlation matrix and LASSO) from the 179 features extracted, the accuracy, sensitivity, specificity, and AUC were 73%, 71%, 71% and 0.74 respectively. This implies that use of radiomics and ML can help to predict treatment response in brain metastasis using pretreatment MRI scan. Future studies will focus on improving prediction accuracy through different radiomics filters and deep learning approaches.

Keywords: Radiomics; machine learning; treatment response; MRI; Gamma Knife

C19

Prediction of Local Recurrence in Head and Neck Cancer Following Chemotherapy Using Radiomic Features from Pre-Radiotherapy MRI

Wun Chin Leong^{1,4}, Hanani Abdul Manan², Caryn Chan Mei Hsien³, Yoke Fui Wong⁴, Erlanianiza Mohd Salim⁵, Noorazrul Yahya¹

¹*Diagnostic Imaging and Radiotherapy, Centre for Diagnostic, Therapeutic and Investigative Sciences (CODTIS), Faculty of Health Science, University of Kebangsaan, Kuala Lumpur, Malaysia*

²*Functional Image Processing Laboratory, Department of Radiology, University of Kebangsaan Malaysia, Kuala Lumpur, Malaysia*

³*Community Health Research Center (ReaCH), Faculty of Health Science, University Kebangsaan Malaysia, Kuala Lumpur, Malaysia*

⁴*Department of Radiotherapy and Oncology, Institut Kanser Negara, Ministry of Health, Putrajaya, Malaysia*

⁵*Department of Radiology, Institut Kanser Negara, Ministry of Health, Putrajaya, Malaysia*

*Correspondence: azrulyahya@ukm.edu.my

ABSTRACT

Head and neck (H&N) cancer treatment is challenging due to critical functions and structures involved. Local recurrence (LC) cause of treatment failure and lead to advanced metastasis. To date, the value of MRI-based radiomics in predicting LC in H&N cancer patients treated with chemotherapy has not been extensively studied. The aim to predict LC in H&N cancer following chemotherapy at pre-RT. A retrospective study enrolled 39 H&N cancer patients treated with CCRT or RT from January 2014 to December 2023. Pre-treatment T1-weighted images, with 228 radiomic features were extracted. Cox proportional hazard model was applied to select radiomic features and clinical characteristics that predict local control (LC). Survival curves were generated using Kaplan-Meier method. The Cox proportional hazards model results indicate a significant relationship between various predictors of diagnosis (HR= 6.86×10^{-9} , 95% CI= 5.39×10^{-9} to 8.73×10^{-9} , $p < 0.01$), squamous cell carcinoma (SCC) types (HR= 38.15, 95% CI= 30.05 to 48.44, $p < 0.01$), tumour (T) (HR= 1.13×10^{-21} , 95% CI= 8.05×10^{-22} to 1.59×10^{-21} , $p < 0.01$), nodule (N) (HR= 2.40×10^{27} , 95% CI= 1.54×10^{27} to 3.72×10^{27} , $p < 0.01$), age (HR= 21.98, 95% CI= 21.05 to 22.95, $p < 0.01$), intensity based (HR= 3.58×10^{56} , 95% CI= 6.35×10^{54} to 2.01×10^{58} , $p < 0.01$), GLCM (HR= 1.88×10^{21} , 95% CI= 1.87×10^{21} to 1.90×10^{21} , $p < 0.01$)/GLRLM (HR= 8.06×10^{260} , 95% CI= 2.89×10^{256} to 2.25×10^{265} , $p < 0.01$)/NGTDM (HR= 0.58, 95% CI= 0.58 to 0.59, $p < 0.01$)/GLSZM (HR= 4.4×10^{17} , 95% CI= 9.77×10^{14} to 1.98×10^{20} , $p < 0.01$) indicates significantly effects on hazard ratio in recurrence risk. Therefore, integrating radiomic features with clinical factors enhance treatment management and achieve better outcomes.

Keywords: Radiomic features; head and neck cancer; Magnetic Resonance Imaging (MRI); chemotherapy; p-radiotherapy

C20

Let's get personal! Molecular Testing for Targeted Therapies in Solid Tumours: HCTM experience

Nurwahyuna R^{1*}, Nur Fa'izah AM², Noor Azlin MH³

¹Department of Pathology, Faculty of Medicine, Universiti Kebangsaan Malaysia

²Department of Radiotherapy and Oncology, Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia

³Department of Diagnostic and Laboratory Services, Hospital Canselor Tuanku Muhriz

*Correspondence: nurwahyuna@ukm.edu.my

ABSTRACT

Personalized medicine has revolutionized the treatment landscape for various cancers. Molecular testing plays a pivotal role in identifying patients who are likely to benefit from targeted therapies. Nevertheless, molecular tests are not readily available in public healthcare settings. This study explores the trends for requests for molecular testing in solid tumours at Hospital Canselor Tuanku Muhriz by analyzing retrospective data from 2020 until 2022. A total of 168 specimens; 124 from primary tumour sites and 44 metastatic sites, underwent molecular testing at external laboratories. There was an upward trend of test requests from 48 in 2020 to 72 in 2022. The most common solid tumours tested were lung (68.5%), colorectal (14.3%), melanoma (5.4%) and lymphoma (4.8%). Other tumours, each comprising less than 2% of the cohort, were gastric, brain, kidney, ovary and endometrium. Lung tumours and melanoma were tested for known driver mutations such as EGFR, ALK, ROS1, BRAF and PDL-1 which are targetable with established medications. Colorectal tumours were tested for KRAS, NRAS, BRAF and MSI status to establish their prognosis and predictive response towards treatment. Testing for PDL-1 expression is a key test often requested to guide treatment in some cancers like non-small cell lung cancer. The findings of this study highlight the increasing demand for various biomarker molecular testing in solid tumours. The transformative potential of biomarker molecular testing in patient care necessitates future efforts to integrate these tests into routine clinical workflows and ensure the affordability of targeted therapies in our setting.

Keywords: Personalized medicine; solid tumours; molecular test; driver mutation

C21

Necrotizing Soft Tissue Infection: Microbiological Distribution from District RegionAmri Dahdi¹, Yazid Bajuri², Norhaslinda Bahaudin³¹Department of Orthopedic and Traumatology, Faculty of Medicine, Universiti Kebangsaan Malaysia²Department of Orthopedic and Traumatology, Faculty of Medicine, Universiti Kebangsaan Malaysia³Department of Orthopedic and Traumatology, Hospital Tuanku Ja'afar, Seremban, N. Sembilan

*Correspondence: P146134@siswa.ukm.edu.my

ABSTRACT

Necrotizing soft tissue infection (NSTI) is fulminant infection with severe systemic inflammatory response and high mortality. Early accurate diagnosis and proper antimicrobials and surgical management are the cornerstones of treatment. NSTI is frequently polymicrobial and the most common organisms are aerobic gram-positive cocci. The objective of this study is to determine microbiology distribution and its sensitivity of NSTI cases from our district cluster center. A retrospective review of all patients (n=110) with documented diagnosis of NSTI together with microbiologic variables of laboratory cultures from August 2020- December 2023 medical records taken from district Hospital of Kuala Pilah and cluster Hospital Jempol and Hospital Tampin were conducted. The results show that the commonest culprit organism involved was *group A Streptococcus* (45%) followed by *Klebsiella sp.* (14%). Based on this data collection, it was proven that the commonest encountered organism of NSTI over Negeri Sembilan district was *Group A Streptococcus*. Treatment wise, recommended for early accurate diagnosis and proper empirical antimicrobials that are effective against aerobic gram-positive cocci and gram-negative rods prior to availability of growth cultures with proper surgical debridement are the cornerstones of treatment.

Keywords: Necrotizing soft tissue infection (NSTI); Negeri Sembilan district; Group A *Streptococcus* (GAS)

C22

Pre-Radiotherapy Balloon Dilatation of the Eustachian Tube: Reducing Radiation-Induced Dysfunction in Nasopharyngeal Carcinoma Patients

Lorna Ting KN^{1,2}, Noor Dina H¹, Lum SG¹, Siti Hajar S², Mohd Razif MY¹, Khairiyah S³

¹Department of Otorhinolaryngology-Head and Neck Surgery, Faculty of Medicine, Universiti Kebangsaan Malaysia

²Department of Otorhinolaryngology-Head and Neck Surgery, Queen Elizabeth Hospital

³Department of Radiotherapy and Oncology, Faculty of Medicine, Universiti Kebangsaan Malaysia

Correspondence: dinahashim@ukm.edu.my

ABSTRACT

Background: Eustachian tube dysfunction (ETD) frequently develops in patients with nasopharyngeal carcinoma (NPC) after radiotherapy, contributing refractory Otitis Media with Effusion (OME). Typically, this condition is dealt with reactively, potentially resulting in complications. Apart from the conventional approach with ventilation tube insertion, balloon dilatation of the Eustachian tube (BDET) offers a safe alternative for treating ETD.

Objective: To evaluate the benefits of BDET prior to radiotherapy in minimizing the risk of ETD among NPC patients.

Methodology: A prospective, randomized controlled trial was conducted among newly diagnosed NPC patients, randomly assigned to intervention group (had Eustachian tube balloon dilatation) or control group (receive no intervention). Assessments included Eustachian Tube Dysfunction Questionnaire-7 (ETDQ-7) scores, tympanometry results, hearing threshold based on pure tone audiometry, with a follow-up of 3 months post-radiotherapy.

Results: 35 patients (70 ears) were initially recruited. Follow-up data at 3 months post-treatment were available for 28 patients (56 ears). No complications observed in the intervention group who underwent BDET under local anesthesia. Post-radiotherapy, significant improvements were documented in ETDQ-7 scores ($p < 0.05$), with greater increases in the intervention group (mean score: 1.45 to 2.15) compared to controls (1.13 to 1.75). Tympanometry findings were similar across groups. In the intervention group, 53.8% showed improved tympanograms, and 73.1% experienced improved hearing thresholds. ABG significantly decreased post-dilatation (mean: 6.35 dB to 5.38 dB), whereas the control group shown an increase of 5 dB.

Conclusion: Balloon Dilatation of the Eustachian Tube (BDET) shows promising effectiveness in reducing radiation-induced Eustachian Tube Dysfunction and related ear complications when performed before radiotherapy. It is suitable for office setting, can be safely integrated into the standard treatment protocol for NPC, ensuring timely commencement of radiotherapy without delays.

Keywords: Nasopharyngeal carcinoma; eustachian tube dysfunction; balloon dilatation of eustachian tube; radiotherapy; otitis media with effusion

C23

Perception of Parents on Adaptive Skills among Preschoolers: A Qualitative StudyMasne K^{1,2*}, Nur Subhanie MA¹, Nor Afifi R^{1,2}, Farahiyah WY^{1,2}¹Occupational Therapy Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia²Centre for Rehabilitation & Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia*Correspondence: masne_kadar@ukm.edu.my**ABSTRACT**

Adaptive skills are practical, everyday skills needed to function and meet the demands of one's environment. These skills are crucial for preschool children, especially during transition to school. While extensive research explores parental perceptions of children with special needs, limited data exists regarding parents' perception of typically developing preschoolers, not to mention their perception on adaptive skills. Therefore, this qualitative study explored the perception of parents on adaptive skills among preschool children between 3 to 6 years old. Fourteen parents of typical preschool children were chosen via purposive sampling and interviewed using a semi-structured layout. Thematic analysis revealed four themes: (1) parents' understanding of adaptive skills; (2) factors influencing the development of adaptive skills; (3) parenting styles and practices and (4) roles and resources in supporting adaptive skills development. The findings showed that parents have a general understanding of their practical implications and emphasized their importance for independence, school readiness, and emotional well-being. They identified numerous factors influencing the development of adaptive skills such as parental roles, environmental factors, child-specific factors and socioeconomic status. Common challenges faced were time constraints, financial limitations, child-related challenges, emotional challenges and challenges in finding resources. These findings underscore the need to develop educational programs to enhance awareness of parents on adaptive skills besides providing support services in addressing specific challenges faced by parents and children. Future research may expand the literature by incorporating diverse cultural and socioeconomic perspectives. The association between parents' perception and child's adaptive skills may also be explored in the future.

Keywords: Parent perception; adaptive skills; preschool children; communication skills; activities of daily living skills; social skills

P1

Understanding Children's Play Perceptions and Play Preferences in Sarawak: A Qualitative Study

Erza Nurfarziany Zalani*, Farahiyah Wan Yunus

Centre for Rehabilitation and Special Needs Studies, Occupational Therapy Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur Campus, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur

*Correspondence: a180864@ukm.edu.my

ABSTRACT

Children engage in play as a form of occupation and as a means of acquiring knowledge about themselves and the world around them. Play can be characterized as a variety of activities independently chosen by children based on their own preferences, enjoyment, and subsequent satisfaction. Play is an essential developmental milestone that contributes in numerous ways to children's development. There are limited studies exploring children's play preferences and experiences. Therefore, this qualitative study aims to investigate the rationale of children's perception of play experiences and play preferences in Sarawak. Ten typical children (five females and five males) aged between 7 and 9 years without any comorbidities participated in the study. An online interview using a semi-structured interview was conducted using Google Meet platform to determine their play preferences and play experiences. Thematic analysis was carried out to analyze and organize the data collected from the interviews. Four main themes emerged which are (i) Playing with Purpose: How Immersive Theatrical Experiences Positively Shape Children's Worlds; (ii) Unlocking the Magic of Play: Exploring the Charismatic Traits of Playful Activities; (iii) Playful Bonds: Unveiling the Charismatic Connections of Playful Interactions and (iv) Where Time, Place, and Weather Unite: Unveiling the Vibrant Worlds of Children's Play Contexts. This study is significant since it offers professionals a deeper understanding of how to work with children. It can also assist occupational therapists and other professionals in incorporating play into their treatment plans, enhancing the use of play in pediatric interventions and evaluations.

Keywords: Semi-structured interview; play; perception; enjoyment; occupational therapy

P2

A Qualitative Study of the Effect of Dialysis on the Performance of Daily Living Activities in Diabetic Clients in the Kuala Selangor Region

Atikah Suib, Farahiyah Wan Yunus

Centre for Rehabilitation and Special Needs Studies, Occupational Therapy Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur Campus, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur

*Correspondence: a182721@ukm.edu.my

ABSTRACT

Dialysis is a well-known treatment for patients with renal failure. Over the years, dialysis patients have increased at an alarming rate. The functional level of dialysis patients is very low compared to non-dialysis diabetes patients. In this study, the researcher aims to determine how dialysis affects diabetes patients' Activity of Daily Living (ADL) performance in Kuala Selangor, as well as which factors contribute to ADL performance. Studies that specifically investigate the ADL performance of dialysis patients with diabetes were found to be limited. This study employed a qualitative method of semi-structured face-to-face interview sessions. Ten hemodialysis participants with stage 3 and stage 4 diabetes, aged thirty and above were recruited. Five main themes emerged from the interview, which are; (i) enhancing everyday life: a fresh perspective on daily activities; (ii) the impact of diabetes on daily living activities; (iii) the impact of dialysis on daily life activities; (iv) functional performance in basic activities of daily living and (v) essential daily life pursuits. Findings from the interview concluded that various types of activities of daily living performance were affected due to low physical body strength among the participants. This study may offer occupational therapy and other rehabilitation disciplines a deeper understanding of the physical capacity of diabetic patients undergoing hemodialysis to perform everyday activities.

Keywords: Diabetes; semi-structured interview; physical activity; activity of daily living; performance

P3

The Quality of Life and Sleep Quality of Health Sciences Undergraduate Students

Noor Azliyana Azmi, Farahiyah Wan Yunus

Centre for Rehabilitation and Special Needs Studies, Occupational Therapy Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur Campus, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur

*Correspondence: a179893@ukm.edu.my

ABSTRACT

The existing literature on the relationship between sleep quality and quality of life seems to be lacking. For health science undergraduate students, having adequate and restful sleep is crucial for the student's optimal physical health, cognitive functioning, and emotional well-being. Students in health science programs often face rigorous academic demands, clinical training, and long hours of study, which can lead to sleep disturbances and poor sleep quality. Poor sleep quality is associated with decreased quality of life. Thus, this study aims to investigate the relationship between sleep quality and the quality of life among health science undergraduate students. A cross-sectional study was conducted using a non-probability convenient sampling technique. A total of 302 participants were selected based on the specified inclusion and exclusion criteria. The survey was distributed through Google Forms using WhatsApp and Facebook. The study utilized the Pittsburgh Sleep Quality Index (PSQI), and the Quality of Life Scale (QOLS). Data was analyzed using the Independent-Samples Mann-Whitney U Test, Independent-Samples Kruskal-Wallis Test, and Spearman's rho. The results reported a strong correlation between sleep quality and the quality of life with $p < 0.001$. Therefore, there is an association between sleep quality and the quality of life among health science undergraduate students. Findings from this study can contribute to the development of strategies and resources aimed at improving sleep habits and managing stress, thereby positively impacting their overall health and quality of life.

Keywords: Cross-sectional study; health sciences; undergraduate students; sleep quality; quality of life

P4

Practice and Perceived Challenges among Clinical Immunologists toward Genetic Testing and Genetic Counseling in Malaysia: A Qualitative Study

Vigneswary Balakrishnan¹, Sharifah Azween Syed Omar¹, Nadiah Sulaiman², Chai-Eng Tan³ and Adli Ali^{1*}

¹*Department of Paediatrics, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, Kuala Lumpur 56000, Malaysia*

²*Centre for Tissue Engineering and Regenerative Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, Kuala Lumpur 56000, Malaysia*

³*Department of Family Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, Kuala Lumpur 56000, Malaysia*

*Correspondence: adli.ali@ppukm.ukm.edu.my

ABSTRACT

Integration of genetic testing and counselling by clinical immunologists are essential in providing definitive diagnosis and holistic care for patients and families with inborn errors of immunity (IEI). However, the practice and perceived challenges of genetic testing and counselling by fellow clinical immunologists in Malaysia remain unclear. Therefore, this study aims to explore the practice and perceived challenges among clinical immunologists toward genetic testing and genetic counselling in Malaysia. This qualitative study employed semi-structured, in-depth one-to-one interviews with clinical immunologists recruited through a purposeful homogeneous sampling method. The findings revealed that clinical immunologists in Malaysia are well-oriented in genetic testing, and they acknowledge the importance of genetic counselling. However, most lack proper genetic counselling in their practice due to challenges underscored in the current study. The emphasis on getting in-house genetic counsellors to provide tailored genetic counselling for patients and families with IEI was a unified calling among all clinical immunologists, as most agreed it would deliver holistic care to families. Discussion on integral gaps in the clinical immunology field in Malaysia was highlighted, and suggestions were made to promote genetic testing and counselling in managing IEI.

Keywords: Practice; perceived challenges; clinical immunologists; genetic testing; genetic counselling; inborn errors of immunity

P5

Reliability, Usability, and Feasibility of the MINAT App among Children Aged 6 to 12 in Malaysia

Wan Sarah Hanan Wan Mohd Hilmi¹, Nor Afifi Razaob^{1,2}, Masne Kadar^{1,2}, Farahiyah Wan Yunus^{1,2}

¹Occupational Therapy Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia.

²Centre for Rehabilitation and Special Needs Studies, Occupational Therapy Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia.

*Correspondence: a183744@ukm.edu.my

ABSTRACT

Self-care, social interaction, education, leisure, and play are important aspects of children's occupational development. This study aimed to determine the reliability, usability, and feasibility of the MINAT application in assessing children's leisure activities. This study used mixed methods: i) quantitative methods to assess reliability, which involved 94 participants; and ii) qualitative methods through interviews among four children aged 6 to 12, their parents, and experienced teachers to evaluate the usability and feasibility of MINAT apps. The data was analyzed using IBM SPSS Version 26 and QDA Miner Lite. The results showed moderate reliability within each domain and high reliability across all domains over time. About 65% of the variance in individual ratings is due to true differences between subjects, with 35% due to errors. For averaged ratings, 89% of the variance is due to true differences between subjects, indicating excellent reliability. The usability and feasibility analysis revealed three main themes: user perceptions and reactions, benefits and personal experiences, and feedback for improvement. The MINAT app is highly reliable, usable, and feasible for assessing children's leisure activities. Recommendations for improvement include adding more accessible content and features to maintain children's engagement.

Keywords: Interests; children; MINAT apps; occupational therapist; leisure activities

P6

Effects of Different Footwear Types and Barefoot on Static and Dynamic Balance among College Students

Nurfarah Najihah, KN, Azira Iqlima, R, Nurhanisah, S, Mohd Murtadha, R, Santibuana, AR*

Faculty of Pharmacy and Health Sciences, Universiti Kuala Lumpur, Royal College of Medicine Perak, 30450 Ipoh, Perak, Malaysia

*Correspondence: santibuana@unikl.edu.my

ABSTRACT

Maintaining proper body balance is essential for preventing falls and injuries. Both static and dynamic balance can be influenced by various types of footwear. However, limited studies have investigated the effects of different footwear types and barefoot on both static and dynamic balance. Additionally, it remains unclear which type of footwear is most effective in enhancing balance among young adults as a preventive measure against falls and injuries. This study aimed to investigate the effects of different footwear types and barefoot on static and dynamic balance among college students. This cross-sectional study involved 48 participants (mean age = 22.42 ± 1.36) selected through purposive sampling. The balance of all participants was assessed under three conditions: wearing sports shoes, wearing flat shoes, and barefoot, using the Balance Error Scoring System Test (BESS) and the Tandem Walk Test (TWT). Data were analyzed using SPSS version 27. The total BESS score indicated significant differences between the sports shoes, flat shoes, and barefoot ($p < 0.001$). Additionally, significant differences were found between footwear types on different surfaces: foam surfaces ($p < 0.01$) and hard surfaces ($p < 0.001$). However, no significant differences in dynamic balance were observed between the footwear and barefoot. In conclusion, different types of footwear and barefoot conditions significantly impact static balance but do not affect dynamic balance. Health promotion campaigns can leverage these findings to educate young adults, especially college students about the importance of appropriate footwear for maintaining balance and reducing the risk of falls and injuries in their daily lives.

Keywords: Footwear; barefoot; static balance; dynamic balance

P7

Validity, Reliability and Profile of Children Handwriting Based on OTulis in Malaysia

Nurnabihah Hashim¹, Nor Afifi Razaob^{1,2}, Masne Kadar^{1,2}, Farahiyah Wan Yunus^{1,2}

¹Occupational Therapy Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia

²Centre for Rehabilitation & Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia

*Correspondence: a181209@siswa.ukm.edu.my

ABSTRACT

Handwriting is essential for children's fine motor skills, cognitive abilities, and literacy. OTulis, a new Bahasa Malaysia handwriting assessment, has shown excellent content validity, but its concurrent validity and reliability needed further evaluation. This study aimed to assess these aspects and profile handwriting skills in Malaysian children aged 4 to 6 using OTulis. A cross-sectional study with purposive sampling included 31 participants (1.71 ± 0.46) for concurrent validity testing with OTulis and the Minnesota Handwriting Assessment (MHA). Reliability and handwriting profiling were assessed in 185 participants (1.59 ± 0.49) using OTulis. Concurrent validity showed a positive correlation between seven OTulis domains and four MHA domains ($r = 0.37-0.99$). No significant correlation was found for the letter recognition domain in OTulis and the rate domain in MHA, with a negative correlation for the fine motor control domain in OTulis and the legibility domain in MHA. Cronbach's alpha for OTulis ranged from 0.89 to 0.90 after removing the visual memory domain. OTulis domain scores (median and IQR) were fine motor control 75 (56-93), fine motor precision 100 (75-100), visual memory 66 (33-100), copy shape 100 (80-100), letter recognition 66 (50-100), uppercase letter formation 35 (65-100), lowercase letter formation 100 (28-100), spacing 0 (0-0), size 100 (66-100), and alignment 100 (0-100). The 0-100 score range of OTulis caters to various skill levels. In conclusion, OTulis shows promise as a valid and reliable tool for assessing handwriting skills beyond fine motor abilities.

Keywords: OTulis; children's handwriting; concurrent validity; internal consistency; profile

P8

Validity and Profile of Children's Activity Preferences of 'MINAT'Wei Shan Tan^{1*}, Nor Afifi Razaob^{1,2}, Masne Kadar^{1,2}, Farahiyah Wan Yunus^{1,2}¹Occupational Therapy Programme, Faculty of Health Sciences, Universiti Kebangsaan Malaysia²Centre for Rehabilitation & Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia

*Correspondence: a179641@siswa.ukm.edu.my

ABSTRACT

Children's participation in various activities is crucial for their growth and quality of life. It is important to use validated assessments and understand children's preferences. 'Penilaian Kecenderungan MINAT', a self-reported assessment measuring children's activity preferences were developed according to the cultural context in Malaysia. This study aimed to determine the validity of 'MINAT', and the profile of children's activity preferences as measured using MINAT among children 6 to 12 years old in Malaysia. A cross-sectional quantitative study with purposive sampling was conducted. MINAT and Preferences for Activity of Children (PAC) were used and involved a total of 218 participants. IBM SPSS Version 26 was used, and the significant level was $p < 0.05$. The validation finding showed that concurrent validity had a moderate to high correlation to PAC (0.53 - 0.79), while the findings of the profile of children showed that there was a significant difference between genders. Physical activity with the mean rank of (male = 125.79, female = 92.91), social (male = 97.05, female = 122.18) and skill-based (male = 97.21, female = 122.02). Similar findings were found in recreational, physical, skill-based, and self-improvement domains, specifically in races. A significant difference in recreational domain was found among children with different household incomes with the mean rank of (<RM 3860 = 54.52, >RM 8319 = 72.02). A profile of MINAT was also obtained in this study. In conclusion, MINAT can be used to measure the activity preferences of children in Malaysia, helping to develop individualized activities to promote participation.

Keywords: MINAT; PAC; children's activity preferences; concurrent validity; profile

P9

MicroRNA Expression Pattern in Complete and Partial Hydatidiform Moles

Wai Kit Chia^{1,2}, Hannah Xin Yi Yeoh¹, Wei -Kang Lee³, Kok Jun Liew³, Chee Choong Hoh³, Teck Yee Khong^{4,5}, Yin Ping Wong^{1*}, Geok Chin Tan^{1*}

¹Department of Pathology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur, Malaysia

²Department of Diagnostic Laboratory Services, Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur, Malaysia

³Codon Genomics S/B, 26, Jalan Dutamas 7, Taman Dutamas Balakong, Selangor, Malaysia

⁴Department of Pathology, SA Pathology, Women's and Children's Hospital, North Adelaide, SA 5006, Australia

⁵University of Adelaide, Adelaide, SA 5000, Australia

*Correspondence: tangc@ppukm.ukm.edu.my and ypwong@ppukm.ukm.edu.my

ABSTRACT

Hydatidiform moles comprised complete (CM) or partial (PM) mole. Although benign, they may progress to choriocarcinoma. MicroRNAs are small, non-coding sequences regulating cellular functions in various tissues, including the placenta. Its role in the pathogenesis of hydatidiform mole remains unclear. This study aimed to determine microRNA expression patterns in CM and PM. Total RNA of CM (n=3), PM (n=3) and non-molar abortus as a control (n=3) were extracted from paraffin embedded tissue blocks. Next-generation sequencing (NGS) and bioinformatics analysis were used to determine microRNA expression. Differentially expressed microRNAs (DEmiRNAs) were identified between the 3 groups using DESeq2. Quantitative real time PCR and digital PCR further validated the expression of these DEmiRNAs. Respective target genes were predicted with TargetScan, TargetMiner and miRDB databases for eleven selected highly DEmiRNAs. NGS data analysis concluded 60 DEmiRNAs in CM and PM, compared to controls. Expression of miR-127-3p, miR-493-5p and miR-106a-5p were downregulated in CM. Expression of miR-584-5p, miR-187-3p and miR-192-5p were upregulated in PM. MicroRNAs downregulated in CM were predicted to share a common target gene *BCCIP*. Conversely, upregulated microRNAs in PM were predicted to share 25 common target genes. Interestingly, *BCCIP*, also known as a BRCA2 and CDKN1A interacting protein, has been found to be involved in genomic stability maintenance, cell cycle regulation and microtubule formation. *BCCIP* deficiency was described in the tumour initiation of breast cancer. Further study on the signalling pathway of CM and PM is required to fully understand the role of microRNAs in the progression of hydatidiform moles.

Keywords: Complete hydatidiform mole; partial hydatidiform mole; molar pregnancy; microRNA; next-generation sequencing

P10

The Effects of Calcium Phosphate Cement Doped with Palm Tocotrienol on Tibial Bone Defect in Ovariectomised Rats through Static Bone Histomorphometry

Siti Sarah MD¹, Wong SK^{1*}, Chin KY¹, Fairus A²

¹Department of Pharmacology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, Kuala Lumpur, 56000, Malaysia

²Department of Anatomy, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, Kuala Lumpur, 56000, Malaysia

*Correspondence: sokkuan@ukm.edu.my

ABSTRACT

Calcium phosphate cement (CPC) is a promising bone substitute material due to its injectability, self-setting properties, moldability, biocompatibility, bioactivity, and osteoconductivity. However, poor osteoinductivity and mechanical properties hinder its clinical use. This study investigated the effects of palm tocotrienol (T3) in addressing the limitations of CPC on bone regeneration in an ovariectomised rat model with tibial bone defects, evaluated through static bone histomorphometry. Twenty-four female Sprague-Dawley rats were split into four groups: sham-operated rats (sham), ovariectomised rats (OVX), ovariectomised rats with bone defect filled with CPC (CPC), and ovariectomised rats with bone defects filled with T3-incorporated CPC (CPC/T3). Eight weeks of treatment were given following a 12 weeks ovariectomy. At the end of the study, the left tibias were harvested and decalcified for two months. Following tissue processing and embedding, tibia samples were stained with haematoxylin and eosin, followed by examination under a microscope. Static bone histomorphometry was measured using the Weibel grid technique. The findings showed that the osteoclast surface of the CPC/T3 group was significantly decreased compared to the sham and OVX groups. Meanwhile, the eroded surface of the CPC/T3 group was significantly increased compared to the OVX group. Osteoblast surface, osteoid surface, and osteoid volume did not show any significant differences between groups. The incorporation of palm T3 into CPC potentially reduces osteoclasts at the bone defect site, which is not observed in those implanted with CPC only.

Keywords: Calcium phosphate cement; tocotrienol; bone defect; osteoporosis

P11

Relationship between Level of Self Satisfaction and Self Confident among Medical Student at National University of Malaysia (UKM) towards Simulation Based Learning

W. Nurfaleana¹, Mohammad Arif Kamarudin¹, Alias Mahmud^{1*}

¹*Department of Medical Education, Faculty of Medicine, Universiti Kebangsaan Malaysia*

*Correspondence: p115392@siswa.ukm.edu.my

ABSTRACT

Simulation-based learning is an essential component of modern medical education in facilitating cognitive and clinical skill development. However, the perception of such methods on students' self-satisfaction and self-confidence has not been extensively studied. This study aims to evaluate the levels of self-satisfaction and self-confidence among Bachelor of Medicine students at Universiti Kebangsaan Malaysia (UKM) and examine the correlation between these variables within the framework of simulation-based learning. A simple random sampling technique was employed to select students from years two to five for participation. The study utilized a pilot-tested questionnaire to ensure reliability, achieving a Cronbach's alpha of 0.99. The data were collected on self-satisfaction and self-confidence and analyzed using descriptive statistics to determine mean scores and standard deviations. The analysis indicated moderate levels of self-satisfaction and self-confidence among students. Self-satisfaction scores ranged from (m = 3.33, SD = 0.53) to (m = 3.53, SD = 0.40) while self-confidence scores exhibited similar values ranging from (m = 3.25, SD = 0.53) to (m = 3.45, SD = 0.44). A strong positive correlation was identified between self-satisfaction and self-confidence with correlation coefficients (r) ranging from 0.77 to 0.85 (p<0.001). The findings suggest that simulation-based learning moderately enhances both self-satisfaction and self-confidence in medical students. These results underline the importance of incorporating simulation-based methods in medical curricula to foster comprehensive skill development in future healthcare professionals.

Keywords: Self-satisfaction; self-confidence; simulation based learning; Bachelor of Medicine students; Universiti Kebangsaan Malaysia

P12

Exploring the Dose-Dependent Impact of Celastrol on Anthropometric Data and Blood Glucose Levels in ApoE-knockout Mice Fed a High-Fat DietArifah, AD¹, Thuhairah Hasrah, AR¹, Vimala, RMTB², Nasibah, A^{1*}¹Department of Medicine, Faculty of Medicine, University Teknologi MARA²Nutrition, Metabolism & Cardiovascular Research Centre, Institute for Medical Research, Setia Alam, Selangor*Correspondence: nasibah@uitm.edu.my**ABSTRACT**

The trend of consuming a hypercaloric (Western) diet in modern society has led to increased body weight and blood glucose levels. Celastrol, an emerging drug derived from *Tripterygium wilfordii* has proven to attenuate these effects in multiple models including, ApoE-knockout mice. However, the outcomes of different celastrol doses remain to be determined. This study aims to investigate the effect of different doses of celastrol on the body weight and blood glucose level of ApoE-knockout mice a fed high-fat diet (HFD). Four groups of male ApoE knockout mice (n=4) were fed with HFD for 12 weeks starting, at 8 weeks of age. During the last 4 weeks, three groups received celastrol intraperitoneally at doses of 1.5 mg, 2 mg, and 2.5 mg/kg/day respectively while the control group received 2% DMSO. Mice's body weight, length, and body mass index (BMI) were recorded weekly and blood glucose was measured at intervals following a 14-hour fast using a glucometer. At the end of the treatment, mice were euthanized using a ketamine-xylazine formulation. Data were analyzed using Kruskal-Wallis and One-way ANOVA tests in GraphPad Prism 10. The findings showed that body weight, BMI, and fasting blood glucose levels initially increased with the HFD, but exhibited a decreasing trend after administering celastrol. Notably, the 2mg/kg dose of celastrol significantly contributed to these observations compared to the control group. This particular dose may be effective for other models and warrants further studies on prolonged Western diet consumption, especially for those dealing with weight gain and high blood glucose levels.

Keywords: Celastrol; ApoE-knockout mice; body weight; BMI; blood glucose

P13

Optimising Caco-2 Cells and *Bifidobacterium Bifidum* Co-Culture to Investigate the Possible Mechanism of Lapatinib-Induced Diarrhoea

Nur Syafina, H¹, Wan Nor I'zzah, WMZ^{1*}, Nur Ain Najiha, R¹, Jesmine, K¹, Narimah, AHH¹ and Hassanain Issam, YT²

¹Department of Biochemistry and Molecular Medicine, Faculty of Medicine, Universiti Teknologi MARA

²Department of Medical Microbiology and Parasitology, Faculty of Medicine, Universiti Teknologi MARA

*Correspondence: wnizzah@uitm.edu.my

ABSTRACT

Lapatinib, an orally administered dual ErbB1/ErbB2 tyrosine kinase inhibitor (TKI), is approved by the FDA for ErbB2-overexpressing breast cancer treatment. However, diarrhoea is an adverse event with an incidence of 58-78% in patients receiving the drug. Previous studies showed a decrease in *Bifidobacterium* sp. in TKI-treated patients with severe diarrhoea, suggesting altered microbial composition, yet the exact mechanism of lapatinib-induced diarrhoea remains vague. This study aims to propose intestinal microflora and its interaction with lapatinib play a role in the mechanism of lapatinib-induced diarrhoea. *Bifidobacterium bifidum* (BB) is selected to investigate if lapatinib inhibits normal intestinal bacteria, leading to diarrhoea. The ability of Caco-2 cells to develop into enterocyte-like phenotypes was adapted to mimic a normal small intestine as an *in vitro* model. The optimum BB concentration that does not affect Caco-2 cell viability in the co-culture model was determined via 3-(4,5-dimethylazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS) assay. Lapatinib inhibition on BB growth co-cultured with Caco-2 was elucidated through an antimicrobial assay. The MTS assay showed 10⁸ CFU/ml of BB provides the optimum Caco-2 cell viability; 86.60±2.73 at 96 hours compared to other BB concentrations. In the antimicrobial assay, lapatinib inhibit BB growth (84.61%±1.68) with an optimum 10⁸ CFU/ml of BB concentration at 96 hours, but no inhibition was observed in the co-culture. In conclusion, a BB concentration of 10⁸ CFU/ml is the most optimal for maintaining Caco-2 cell viability and growth, at 96 hours, for co-culture development. Nevertheless, possible mechanism of lapatinib-induced diarrhoea via inhibition of normal intestinal bacteria growth requires further observation.

Keywords: Lapatinib; Caco-2; *Bifidobacterium bifidum*; MTS assay; antimicrobial assay

P14

Ultrastructural Characteristics of Placentas in Pregnancies Complicated with Fetal Growth Restriction

Che En Tan¹, Kevin Boon Ping Ku¹, Wirda Indah Farouk², Rahana Abd Rahman³, Geok Chin Tan^{1*}, Yin Ping Wong^{1*}

¹Department of Pathology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur, Malaysia

²Department of Diagnostic Laboratory Services, Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak

³Department of Obstetrics and Gynaecology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur, Malaysia

*Correspondence: ypwong@ppukm.ukm.edu.my and tangc@ppukm.ukm.edu.my

ABSTRACT

Fetal growth restriction (FGR) is a serious pregnancy complication associated with higher perinatal morbidity and mortality, primarily related to abnormalities in placental anatomical structure or function. Even now, the exact causes of FGR remain poorly understood. This study aimed to identify the ultrastructural features of placentas in pregnancies complicated with FGR. Control placentas (n=11) and FGR placenta samples (n=10) were collected from consented mothers. Placental tissues were processed in accordance with the manufacturer's guide and analysed with transmission electron microscopy (TEM). Parameters such as the length and width of microvilli, the numbers of microvilli per 1 micron, basement membrane thickness and diffusion distance between the two groups were measured and recorded. A significant increase in the diffusion distance was observed in the villi of FGR placentas ($289.90 \pm 44.40\mu\text{m}$) compared to those of normal ($215.60 \pm 57.54\mu\text{m}$) ($p = 0.0019$). There were no significant differences in the microvilli length, width, number of microvilli per 1 micron or basement membrane thickness between the control and FGR groups ($p > 0.05$). The increase in the diffusion distance from maternal blood to fetal vessels in placentas complicated with FGR may impede effective transfer of oxygen supply and necessary nutrients to the fetus through the placenta, resulting in compromised fetal growth and development. Further studies with a larger sample size are however needed to elucidate this association.

Keywords: Fetal growth restriction; placenta; transmission electron microscopy (tem); ultrastructure; diffusion distance

P16

Mapping Regorafenib's Protein Targets in Colorectal Cancer: Insights from Molecular Fingerprinting

Farina Syazwani Md Azhan, Muhammad Redha Abdullah Zawawi, Teck Yew Low, Pey Yee Lee*

UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur

*Correspondence: leepeyyee@ukm.edu.my

ABSTRACT

Globally, colorectal cancer (CRC) ranks third in incidence and second in cancer-related mortality. Cytotoxic chemotherapy is the standard first- and second-line treatment for metastatic CRC (mCRC) patients, either alone or in combination with targeted agents. Unfortunately, most mCRC patients experience disease progression after standard treatments and have limited subsequent therapeutic options. Regorafenib (REG), an FDA-approved multi-kinase inhibitor, has demonstrated efficacy in mCRC patients who are refractory to all previous therapies. Recent evidences suggest that combining REG with other targeted agents may improve outcomes, potentially moving its use to earlier treatment lines. However, the complete drug targets and molecular mechanism of REG remain unclear. In this work, we aim to elucidate the drug target profile of REG in CRC, gaining insights into the drug's mechanism of action. To achieve this, we computed the structural similarity of REG with FDA-approved drugs to infer drug targets from similar drug-ligand interactions by using molecular fingerprinting technique. Using the Tanimoto coefficient (cut off value > 0.7), we found nine drugs with a likely similar structure to REG. Based on molecular fingerprinting results, we discovered ten new potential proteins targets of REG that have not been previously reported. Notably, the Cystic Fibrosis Transmembrane Conductance Regulator (CFTR), whose mutations can lead to chronic inflammation in the intestines, is one of the new potential targets identified. By identifying new protein targets of REG computationally, we aim to create a comprehensive map of REG's protein targets within CRC, potentially refining treatment strategies and improving therapeutic outcomes for mCRC patients.

Keywords: Colorectal cancer; regorafenib; targeted therapy; molecular fingerprinting; kinase inhibitor

P17

Enrichment of GBP7 Mutation in Patients with Primary Aldosteronism

Fatin Athirah Pauzi¹, Nurfatim Syazreen Mod Rizam¹, Munkit Choy², Simon Williams², Muaatamarulain Mustangin³, Geok Chin Tan³, Xilin Wu⁴, Elisabeth Ng^{4,5}, Kate Laycock⁴, Azraai Bahari Nasruddin⁶, Norlela Sukor^{1,7}, Morris Jonathan Brown⁴, Bernard Keaveney², Elena Aisha Azizan^{1*}

¹Department of Medicine, Faculty of Medicine, The National University of Malaysia (UKM) Medical Centre, Malaysia

²Division of Cardiovascular Sciences, University of Manchester, United Kingdom

³Department of Pathology, Faculty of Medicine, The National University of Malaysia (UKM) Medical Centre, Malaysia

⁴Department of Clinical Pharmacology and Precision Medicine, William Harvey Research Institute, Queen Mary University of London, United Kingdom

⁵Centre for Endocrinology & Metabolism, Hudson Institute of Medical Research, Victoria, Australia

⁶Department of Endocrinology, Hospital Putrajaya, Malaysia

⁷Hospital Canselor Tuanku Muhriz Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur, Malaysia

*Correspondence: elena.azizan@ukm.edu.my

ABSTRACT

Primary aldosteronism (PA), showing excess production of the hormone aldosterone from the adrenal glands, is probably the commonest curable cause of hypertension. However, genome-wide association studies have only identified ~3.5% of the genetic contribution in blood pressure with most hits occurring in the non-coding regions. Whole genome sequencing (WGS), a powerful technology to detect all the genetic sequences in an individual, were utilized to identify the heritable mutations associated with PA. Based on the WGS results of 134 subjects, an enrichment of rare missense mutations in GBP7 was found in PA patients of Black ethnicity or in PA patients who have small aldosterone-producing micronodules (APMs). GBP7 is of interest as it encodes for a guanylate-binding protein that can hydrolyze GTP to both GDP and GMP, a process that plays a role in aldosterone synthesis pathway. The mutations in GBP7 has been verified to be true by Sanger Sequencing and overexpression of GBP7 in HAC15 cells significantly increased expression of CYP11B2 mRNA, the aldosterone synthase gene, by 1.57 ± 0.21 fold ($p=0.001$). This was similar to the increase in CYP11B2 mRNA expression seen with angiotensin II-stimulated cells. Whereas silencing of GBP7 in HAC15 cells, significantly reduced CYP11B2 expression. These results suggest that alteration in the expression of GBP7 affects aldosterone production highlighting that GBP7 may play a potential role in PA pathogenesis especially those with idiopathic hyperaldosteronism and APMs.

Keywords: Primary aldosteronism; aldosterone; CYP11B2; GBP7; whole genome sequencing

P18

Exploring NPFS Mutations in Pediatric Patients with Congenital and Steroid-Resistant Nephrotic Syndrome

Jun Xin Lee, Noor Akmal Shareela binti Ismail*

Department of Biochemistry, Faculty of Medicine, Universiti Kebangsaan Malaysia

*Correspondence: nasismail@ukm.edu.my

NPFS1 and NPFS2 are both gene components of kidney that encodes for nephrin and podocin respectively. They play a role in progression of congenital (CNS) and steroid resistant (SRNS) nephrotic syndrome. Hence, this study aims to determine the prevalence and renal outcome of NPFS mutations among CNS and SRNS pediatric patients. We also aim to identify potential predictors of NPFS mutations in this patient cohort. Overall, there were 33 studies involving 2123 patients being screened for NPFS1, whereas 2889 patients from 40 studies were screened for NPFS2 mutations. Patients' mean age was 4.9 ± 1 years (ranged from birth – 18 years) and 56% was male (n=1281). Using random effects model, the pooled proportion of NPFS1 mutations among CNS and SRNS pediatric patients was 0.15 (95% CI 0.09; 0.24, $p < 0.001$, $I^2 = 92.0\%$). The pooled proportion of NPFS2 mutations was slightly lower 0.11 (95% CI 0.08; 0.14, $p < 0.001$, $I^2 = 73.8\%$). Among the 18 studies that reported on ESRF, the pooled proportion was 0.47 (95% CI 0.34; 0.61, $p < 0.001$, $I^2 = 75.4\%$). Our study showed that both NPFS1 ($\text{OR} = 1.16$, $p = 0.35$) and NPFS2 ($\text{OR} = 5.49$, $p = 0.08$) mutations did not predict ESRF in CNS and SRNS pediatric patients. Nevertheless, NPFS2 mutation from the European continents had a significantly higher risk of getting ESRF ($p < 0.05$, $\text{OR} = 1.3$, $\text{OR} = 7.97$, 95% CI 0.30; 2.30) but not in NPFS1 mutation. We recommend NPFS mutations' screening for earlier diagnosis and avoid unnecessary steroid treatments. More data are needed to better understand the impact of NPFS mutations among pediatric CNS and SRNS patients.

Keywords: NPFS1; NPFS2; congenital nephrotic syndrome; steroid resistant nephrotic syndrome; pediatric; end stage renal failure

P19

Effects of Kelulut Honey on the Frontal Lobe and Cerebellum in Amyloid β -induced Alzheimer's Disease Rat ModelSuhilen TS, Shaik A, Teh SL, Yahaya MF**Department of Anatomy, Faculty of Medicine, Universiti Kebangsaan Malaysia**Correspondence: mfairuzy@ukm.edu.my**ABSTRACT**

Alzheimer's disease (AD) is a progressive neurodegenerative disorder and is marked by cognitive decline and memory loss, due to β -amyloid plaques and neurofibrillary tangles in the brain. Kelulut honey, known for its antioxidant and neuroprotective properties, was tested for its effects on AD-induced rat models, focusing on the frontal lobe and cerebellum. Twenty-six male Sprague-Dawley rats, weighing 280-380g, were randomly divided into three groups: control, disease (AB), and treatment with Kelulut honey (KH). The AB and KH groups underwent stereotaxic surgery, receiving intrahippocampal injection of 2.5 $\mu\text{g}/\mu\text{l}$ of $\text{A}\beta_{1-42}$. One-week post-surgery, the treatment group received Kelulut honey at a dose of 1g/kg, for four weeks. The rats were then sacrificed and their brains were processed for histochemical staining using H&E and TUNEL assay, and ELISA. Findings indicate that Kelulut honey helps maintain the number of pyramidal cells in the frontal lobe ($p < 0.05$), which would otherwise decrease in AD. The TUNEL assay showed no significant difference in apoptotic cells between the groups in both the cerebellum and the frontal lobe. However, ELISA for SOD showed significant differences between AB and control group ($p < 0.05$) in the cerebellum, and treatment with KH significantly elevates ($p < 0.05$) the SOD level. We also found no significant differences between all groups for $\text{A}\beta_{42}$, p-tau, MDA and NF κ B. Our findings indicate that Kelulut honey treatment was able to reduce amyloid plaque deposition and elevated the antioxidant levels, suggesting its potential to alleviate AD-related pathology.

Keywords: Intrahippocampal injection; β -amyloid; tau protein; antioxidant

P20

Proteomic profiling reveals interferon pathway upregulation in FOLFOX-resistant colorectal cancer cells

Yeoh Yeelon, Low Teck Yew, Nadiah Abu, Lee Pey Yee*

*UKM Medical Molecular Biology Institute (UMBI), Jalan Yaacob Latif, Bandar Tun Razak, 56000 Cheras,
Wilayah Persekutuan Kuala Lumpur*

*Correspondence: leepeyyee@ukm.edu.my

ABSTRACT

Colorectal cancer (CRC) patients receiving FOLFOX chemotherapy may eventually develop drug resistance, which remains poorly understood. To elucidate the mechanisms behind FOLFOX resistance, we generated FOLFOX-resistant CRC cell lines by exposing HCT-116 cells to FOLFOX at clinically relevant doses for ten cycles. The development of drug resistance was validated with cell viability assays (MTT). The cell lysates of both resistant and parental cells (before and after 24 hours of FOLFOX treatment) were then collected for mass spectrometry-based proteomics and bioinformatic analysis. As a result, we identified 9,477 proteins, of which 8,388 were quantified. Parental cells treated with FOLFOX showed significant overexpression of proteins associated with DNA damage response and cell cycle arrest (TP53, CDKN1A, CCNE1 and CCNA2). On the contrary, resistant cells did not exhibit enrichment of such pathways. Such observations are consistent with FOLFOX being a DNA-damaging agent and suggest a mechanism for the development of resistance. A comparison of parental and resistant cells before treatment showed that 51 out of 87 identified interferon-stimulated genes (ISGs) were significantly upregulated in the resistant cells. After 24 hours of FOLFOX treatment, there were 31 upregulated ISGs in the resistant cells, indicating the probable involvement of interferon pathways in FOLFOX resistance. Notable examples of these upregulated ISGs include ISG15, IFI35, IFIH1, OAS3, and STAT1. Our data highlights ISGs as potential biomarkers for stratifying responsive and resistant patients to FOLFOX treatment. Further validation in clinical settings is needed to establish their predictive utility and potential translation into clinical practice.

Keywords: Colorectal cancer; FOLFOX; interferon pathways; interferon-stimulated genes; proteomics

P21

Precision Medicine Unveils the Molecular Profiles of Extracellular Matrix Anabolism, Inflammation, and Nociception Tailored to the Severity of Degenerative Disc Disease in Humans

Nurul Fariha Z^{1,2}, Izzat Z², Kieran J^{2,3}, Karina D⁴, Aoibhinn K⁴, Ahmad Hafiz K⁵, Muhamad Nazrin M⁵, Sabarul Afian M⁵, Elvy Suhana M R¹, Fairus A¹, David P F^{2,4}, Abhay P², Aiden D³, Isma Liza M I^{1,2,4}

¹Department of Anatomy, Faculty of Medicine, Universiti Kebangsaan Malaysia, ²CÚRAM, SFI Research Centre for Medical Devices, University of Galway, Ireland, ³Orthopedic Surgery, University Hospital Galway, Ireland, ⁴Pharmacology and Therapeutics, School of Medicine, University of Galway, Ireland, ⁵Department of Orthopaedics and Traumatology, Faculty of Medicine, Universiti Kebangsaan Malaysia

*Correspondence: ismaliza.mohdisa@ukm.edu.my

ABSTRACT

Intervertebral disc (IVD) degeneration is one of the primary etiological factors contributing to low back pain (LBP). It is mediated by an imbalance of extracellular matrix (ECM) homeostasis and increased inflammation that induces innervation, causing nociception. We hypothesised that human degenerative IVD differentially expressed molecular markers of ECM anabolism, inflammation and nociception tailored to the severity of the degenerative disc disease (DDD) using precision medicine. We adopted a multicentre prospective study of transcriptomic profiles in IVD tissues collected from DDD patients with LBP and healthy IVD from idiopathic scoliosis patients who underwent spinal surgery (Merlin Park Hospital and University Hospital Galway) and body donors (Silent Mentor Programme, Malaysia). We graded the patient's MRI images using *Pfirrmann* grading and the donor's IVD histology using *Thompson* grading. We conducted RNA sequencing (Cambridge, UK) for transcriptome, bioinformatics and marker validation by RT-qPCR. We determined DDD patients with *Pfirrmann* grades IV ($n=11$) and grades V ($n=11$). There was an increased age (range 40 to 70 years) as the severity of DDD advanced; however, there were no gender differences. The control group procurement, $n=8$ for adult healthy IVD and $n=4$ for adolescent healthy IVD. Transcriptome revealed differentially expressed genes of ECM anabolic (COL1A2 and COL2A1), IL-1 β and pro-nociceptive NGF and TRPV4 in DDD. We confirmed an increased trend of COL1A2 expression and reduced trend for COL2A1 expression as the severity of DDD advanced, respectively. These findings indicate dysregulation of ECM anabolism and upregulation of inflammation and nociception, offering insight into the molecular mechanism underlying the severity of DDD, potentially imply the therapeutic targets for treating LBP through precision medicine.

Keywords: Low back pain; intervertebral disc degeneration; transcriptome; inflammation; nociception. extracellular matrix



FULL PROGRAM

Scan Me



<https://tinyurl.com/ProgramRW2024>